

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US05/007106

International filing date: 04 March 2005 (04.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/551,440
Filing date: 09 March 2004 (09.03.2004)

Date of receipt at the International Bureau: 07 April 2005 (07.04.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse

1302474

THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office

March 31, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE.

APPLICATION NUMBER: 60/551,440

FILING DATE: *March 09, 2004*

RELATED PCT APPLICATION NUMBER: *PCT/US05/07106*



Certified by

Under Secretary of Commerce
for Intellectual Property
and Director of the United States
Patent and Trademark Office

23026 U.S. PTO
030904

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a **PROVISIONAL APPLICATION FOR PATENT** under 37 CFR 1.53 (c).

22264 U.S. PTO
60/551440
030904

		DOCKET NUMBER		21548PV	
INVENTOR(S)					
Given Name (first and middle [if any])		Family Name or Surname		Residence (City and either State or Foreign Country)	
Matthew M. Peter D. John S. Thorsten E. Terry A.		Morrisette Williams Wai Fisher Lyle		Wayne, PA Harleysville, PA Harleysville, PA Hatfield, PA Lederach, PA	
<input type="checkbox"/> Additional inventors are being named on the separately numbered sheets attached hereto					
TITLE OF THE INVENTION (500 characters max)					
HIV INTEGRASE INHIBITORS					
CORRESPONDENCE ADDRESS					
Direct all Correspondence to: Merck & Co., Inc. Patent Department - RY60-30 P.O. Box 2000 Rahway					
STATE		New Jersey		ZIP CODE	
				07065	
COUNTRY		U.S.A.		Customer Number	
				000210	
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification		Number of Pages		62	
<input type="checkbox"/> Drawing(s)		Number of Sheets		<input type="checkbox"/> CD(s), Number	
<input type="checkbox"/> Application Data Sheet. See 37 CFR 1.76				<input type="checkbox"/> Other (specify)	
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT (check one)					
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees				FILING FEE AMOUNT (\$)	
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: 13-2755				\$160.00	

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☒ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are: _____

Respectfully submitted,

SIGNATURE Kenneth R. Walton

Date 03/09/2004

TYPED or PRINTED NAME Kenneth R. Walton

REGISTRATION NO. 32,951

TELEPHONE 732 594 3462

(if appropriate)

NOTE: Mail to Mail Stop Provisional Application

EXPRESS MAIL CERTIFICATE	
DATE OF DEPOSIT	<u>March 9, 2004</u>
EXPRESS MAIL NO.	<u>EL 989589934 US</u>
I HEREBY CERTIFY THAT THIS CORRESPONDENCE IS BEING DEPOSITED WITH THE UNITED STATES POSTAL SERVICE AS EXPRESS MAIL "POST OFFICE TO ADDRESSEE" ON THE ABOVE DATE IN AN ENVELOPE ADDRESSED TO COMMISSIONER FOR PATENTS, P.O. BOX 1450, ALEXANDRIA, VA 22313-1450.	
MAILED BY	<u>Kenneth R. Walton</u>
DATE	<u>March 9, 2004</u>

In Duplicate

TITLE OF THE INVENTION
HIV INTEGRASE INHIBITORS

FIELD OF THE INVENTION

5 The present invention is directed to hydroxy tetrahydronaphthyrine dione and hydroxy hexahydronaphthyrine dione compounds and pharmaceutically acceptable salts thereof, their synthesis, and their use as inhibitors of the HIV integrase enzyme. The compounds and pharmaceutically acceptable salts thereof of the present invention are useful for preventing or treating infection by HIV and for preventing or treating or delaying the onset of AIDS.

10

BACKGROUND OF THE INVENTION

 A retrovirus designated human immunodeficiency virus (HIV) is the etiological agent of the complex disease that includes progressive destruction of the immune system (acquired immune deficiency syndrome; AIDS) and degeneration of the central and peripheral nervous system. This virus was previously known as LAV, HTLV-III, or ARV. A common feature of retrovirus replication is the insertion by virally-encoded integrase of proviral DNA into the host cell genome, a required step in HIV replication in human T-lymphoid and monocytoid cells. Integration is believed to be mediated by integrase in three steps: assembly of a stable nucleoprotein complex with viral DNA sequences; cleavage of two nucleotides from the 3' termini of the linear proviral DNA; covalent joining of the recessed 3' OH termini of the proviral DNA at a staggered cut made at the host target site. The fourth step in the process, repair synthesis of the resultant gap, may be accomplished by cellular enzymes.

 Nucleotide sequencing of HIV shows the presence of a pol gene in one open reading frame [Ratner, L. et al., Nature, 313, 277(1985)]. Amino acid sequence homology provides evidence that the pol sequence encodes reverse transcriptase, integrase and an HIV protease [Toh, H. et al., EMBO J. 4, 1267 (1985); Power, M.D. et al., Science, 231, 1567 (1986); Pearl, L.H. et al., Nature, 329, 351 (1987)]. All three enzymes have been shown to be essential for the replication of HIV.

 It is known that some antiviral compounds which act as inhibitors of HIV replication are effective agents in the treatment of AIDS and similar diseases, including reverse transcriptase inhibitors such as azidothymidine (AZT) and efavirenz and protease inhibitors such as indinavir and nelfinavir. The compounds of this invention are inhibitors of HIV integrase and inhibitors of HIV replication. The inhibition of integrase in vitro and HIV replication in cells is a direct result of inhibiting the strand transfer reaction catalyzed by the recombinant integrase in vitro in HIV infected

cells. The particular advantage of the present invention is highly specific inhibition of HIV integrase and HIV replication.

The following references are of interest as background:

US 6380249, US 6306891, and US 6262055 disclose 2,4-dioxobutyric acids and acid esters useful as HIV integrase inhibitors.

WO 01/00578 discloses 1-(aromatic- or heteroaromatic-substituted)-3-(heteroaromatic substituted)-1,3-propanediones useful as HIV integrase inhibitors.

US 2003/0055071 (corresponding to WO 02/30930), WO 02/30426, and WO 02/55079 each disclose certain 8-hydroxy-1,6-naphthyridine-7-carboxamides as HIV integrase inhibitors.

WO 02/036734 discloses certain aza- and polyaza-naphthalenyl ketones to be HIV integrase inhibitors.

WO 03/016275 discloses certain compounds having integrase inhibitory activity.


WO 03/35076 discloses certain 5,6-dihydroxypyrimidine-4-carboxamides as HIV integrase inhibitors, and WO 03/35077 discloses certain N-substituted 5-hydroxy-6-oxo-1,6-dihydropyrimidine-4-carboxamides as HIV integrase inhibitors.

WO 03/062204 discloses certain hydroxynaphthyridinone carboxamides that are useful as HIV integrase inhibitors.

WO 04/004657 discloses certain hydroxypyrrole derivatives that are HIV integrase inhibitors.

SUMMARY OF THE INVENTION

The present invention is directed to hydroxy polyhydronaphthyridine dione compounds. These compounds are useful in the inhibition of HIV integrase, the prevention of infection by HIV, the treatment of infection by HIV and in the prevention, treatment, and delay in the onset of AIDS and/or ARC, either as compounds or their pharmaceutically acceptable salts or hydrates (when appropriate), or as pharmaceutical composition ingredients, whether or not in combination with other HIV/AIDS antivirals, anti-infectives, immunomodulators, antibiotics or vaccines. More particularly, the present invention includes compounds of Formula I, and pharmaceutically acceptable salts thereof:



wherein:

R^I is $-C_{1-6}$ alkyl substituted with R^J , wherein R^J is:

- (A) aryl or aryl fused to a 5- or 6-membered heteroaromatic ring containing from 1 to 4 heteroatoms independently selected from N, O and S, wherein the aryl or fused aryl is
- (i) optionally substituted with from 1 to 5 substituents each of which is independently:
- (1) $-C_{1-6}$ alkyl, which is optionally substituted with $-OH$, $-O-C_{1-6}$ alkyl, $-O-C_{1-6}$ haloalkyl, $-CN$, $-NO_2$, $-N(R^A)R^B$, $-C(=O)N(R^A)R^B$, $-C(=O)RA$, $-CO_2RA$, $-S(O)_nRA$, $-SO_2N(R^A)R^B$, $-N(R^A)C(=O)R^B$, $-N(R^A)CO_2R^B$, $-N(R^A)SO_2R^B$, $-N(R^A)SO_2N(R^A)R^B$, $-OC(=O)N(R^A)R^B$, or $-N(R^A)C(=O)N(R^A)R^B$,
 - (2) $-O-C_{1-6}$ alkyl,
 - (3) $-C_{1-6}$ haloalkyl,
 - (4) $-O-C_{1-6}$ haloalkyl,
 - (5) $-OH$,
 - (6) halo,
 - (7) $-CN$,
 - (8) $-NO_2$,
 - (9) $-N(R^A)R^B$,
 - (10) $-C(=O)N(R^A)R^B$,
 - (11) $-C(=O)RA$,
 - (12) $-CO_2RA$,
 - (13) $-SRA$,
 - (14) $-S(=O)RA$,
 - (15) $-SO_2RA$,
 - (16) $-SO_2N(R^A)R^B$,
 - (17) $-N(R^A)SO_2R^B$,
 - (18) $-N(R^A)SO_2N(R^A)R^B$,
 - (19) $-N(R^A)C(=O)R^B$,
 - (20) $-N(R^A)C(=O)-C(=O)N(R^A)R^B$, or
 - (21) $-N(R^A)CO_2R^B$, and
- (ii) optionally substituted with 1 or 2 substituents each of which is independently:
- (1) aryl,

- (2) -C₁₋₆ alkyl substituted with aryl,
- (3) -HetA,
- (4) -C(=O)-HetA; or
- (5) -HetB;

wherein each HetA is independently a C₄₋₇ azacycloalkyl or a C₃₋₆ diazacycloalkyl, either of which is optionally substituted with from 1 to 3 substituents each of which is independently oxo or C₁₋₆ alkyl; and

wherein each HetB is a 5- or 6-membered heteroaromatic ring containing from 1 to 4 heteroatoms independently selected from N, O and S, wherein the heteroaromatic ring is optionally substituted with from 1 to 4 substituents each of which is independently halo, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, or hydroxy; or

- (B) a 5- or 6-membered heteroaromatic ring containing from 1 to 4 heteroatoms independently selected from N, O and S; wherein the heteroaromatic ring is:

- (i) optionally substituted with from 1 to 4 substituents each of which is independently halogen, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, or hydroxy, and
- (ii) optionally substituted with 1 or 2 substituents each of which is independently aryl or -C₁₋₆ alkyl substituted with aryl;

R², R³, R⁴ and R⁵ are defined as follows:

- (A) R², R³, R⁴ and R⁵ are each independently:

- (1) -H,
- (2) -C₁₋₆ alkyl, which is optionally substituted with -OH, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, -CN, -N(RA)RB, -C(=O)N(RA)RB, -C(=O)RA, -CO₂RA, -S(O)_nRA, -SO₂N(RA)RB, -N(RA)C(=O)RB, -N(RA)CO₂RB, -N(RA)SO₂RB, -N(RA)SO₂N(RA)RB, -N(RA)C(=O)N(RA)RB, or -OC(=O)N(RA)RB,
- (3) -C₁₋₆ haloalkyl,
- (4) CycA,
- (5) AryA,
- (6) HetC, or
- (7) -C₁₋₆ alkyl substituted with CycA, AryA, or HetC;

- (B) R² and R⁴ together with the carbon atoms to which each is attached form a carbon-carbon double bond; and R³ and R⁵ are each independently as defined in part A above;

(C) R² and R³ together with the carbon atom to which they are both attached form a 3- to 8-membered saturated carbocyclic ring which is optionally substituted with from 1 to 4 substituents each of which is independently -OH, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, or -O-C₁₋₆ haloalkyl; and R⁴ and R⁵ are each independently as defined in part A above; or

(D) R⁴ and R⁵ together with the carbon atom to which they are both attached form a 3- to 8-membered saturated carbocyclic ring which is optionally substituted with from 1 to 4 substituents each of which is independently -OH, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, or -O-C₁₋₆ haloalkyl; and R² and R³ are each independently as defined in part A above;

R⁶ is:

- (1) -H,
- (2) -C₁₋₆ alkyl, which is optionally substituted with -OH, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, -CN, -N(RA)RB, -C(=O)N(RA)RB, -C(=O)RA, -CO₂RA, -S(O)_nRA, -SO₂N(RA)RB, -N(RA)C(=O)RB, -N(RA)CO₂RB, -N(RA)SO₂RB, -N(RA)SO₂N(RA)RB, -N(RA)C(=O)N(RA)RB, or -OC(=O)N(RA)RB,
- (3) -C₁₋₆ haloalkyl,
- (4) CycA,
- (5) AryA,
- (6) HetC, or
- (7) -C₁₋₆ alkyl substituted with CycA, AryA, or HetC;

R⁷ and R⁸ are each independently:

- (1) -H,
- (2) -C₁₋₆ alkyl, which is optionally substituted with -OH, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, -CN, -N(RA)RB, -C(=O)N(RA)RB, -C(=O)RA, -CO₂RA, -S(O)_nRA, -SO₂N(RA)RB, -N(RA)C(=O)RB, -N(RA)CO₂RB, -N(RA)SO₂RB, -N(RA)SO₂N(RA)RB, -N(RA)C(=O)N(RA)RB, or -OC(=O)N(RA)RB,
- (3) -C₁₋₆ haloalkyl,
- (4) -C(=O)RA,
- (5) -CO₂RA,
- (6) -C(=O)N(RA)RB,
- (7) -N(RA)SO₂N(RA)RB,

- (8) -RK,
 (9) -C(=O)-RK,
 (10) -C(=O)N(RA)-RK,
 (11) -C(=O)N(RA)-C₁₋₆ alkylene-RK, or
 5 (12) -C₁₋₆ alkyl substituted with -RK, -C(=O)-RK, -C(=O)N(RA)-RK, or -C(=O)N(RA)-C₁₋₆ alkylene-RK;

or alternatively R⁷ and R⁸ together with the carbon atom to which they are both attached form a 3- to 8-membered saturated carbocyclic ring which is optionally substituted with from 1 to 4 substituents each of
 10 which is independently halogen, -OH, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, or -O-C₁₋₆ haloalkyl;

R⁹ is:

- (1) -H,
 (2) -C₁₋₆ alkyl, which is optionally substituted with -OH, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl,
 15 -CN, -N(RA)RB, -C(=O)N(RA)RB, -C(=O)RA, -CO₂RA, -S(O)_nRA, -SO₂N(RA)RB,
 -N(RA)C(=O)RB, -N(RA)CO₂RB, -N(RA)SO₂RB, -N(RA)SO₂N(RA)RB,
 -N(RA)C(=O)N(RA)RB, or -OC(=O)N(RA)RB,
 (3) -C₁₋₆ haloalkyl,
 (4) CycA,
 20 (5) AryA,
 (6) HetC, or
 (7) -C₁₋₆ alkyl substituted with CycA, AryA, or HetC;

each n is independently an integer equal to zero, 1, or 2;

25 each R^A is independently H or C₁₋₆ alkyl;

each R^B is independently H or C₁₋₆ alkyl;

30 each R^K is independently CycA, AryA, or HetC;

each CycA is independently a C₃₋₈ cycloalkyl, which is optionally substituted with from 1 to 4 substituents each of which is halogen, -OH, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, or -O-C₁₋₆ haloalkyl;

each AryA is independently an aryl, which is

- 5 (a) optionally substituted with from 1 to 5 substituents each of which is independently -C₁₋₆ alkyl, -C₁₋₆ alkylene-OH, -C₁₋₆ alkylene-O-C₁₋₆ alkyl, -C₁₋₆ alkylene-O-C₁₋₆ haloalkyl, -C₁₋₆ alkylene-N(RA)R^B, -C₁₋₆ alkylene-C(=O)N(RA)R^B, -C₁₋₆ alkylene-C(=O)RA, -C₁₋₆ alkylene-CO₂RA, -C₁₋₆ alkylene-S(O)_nRA, -O-C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ haloalkyl, -OH, halo, -N(RA)R^B, -C(=O)N(RA)R^B, -C(=O)RA, -CO₂RA, -S(O)_nRA, or -SO₂N(RA)R^B, and
- 10 (b) optionally substituted with C₃₋₈ cycloalkyl, aryl, HetD, or -C₁₋₆ alkyl substituted with C₃₋₈ cycloalkyl, aryl, or HetD;

each HetC is independently a 4- to 7-membered saturated or unsaturated heterocyclic ring containing at least one carbon atom and from 1 to 4 heteroatoms independently selected from N, O and S, wherein the heterocyclic ring is

- 15 (a) optionally substituted with from 1 to 4 substituents each of which is halogen, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, OH, or oxo, and
- (b) optionally substituted with C₃₋₈ cycloalkyl, aryl, HetD, or -C₁₋₆ alkyl substituted with C₃₋₈ cycloalkyl, aryl, or HetD;

- 20 each HetD is independently a 4- to 7-membered saturated or unsaturated heterocyclic ring containing at least one carbon atom and from 1 to 4 heteroatoms independently selected from N, O and S; and

each aryl is independently (i) phenyl or (ii) a 9- or 10-membered bicyclic, fused carbocyclic ring system in which at least one ring is aromatic.

- 25 The present invention also includes pharmaceutical compositions containing a compound of the present invention and methods of preparing such pharmaceutical compositions. The present invention further includes methods of treating AIDS, methods of delaying the onset of AIDS, methods of preventing AIDS, methods of preventing infection by HIV, and methods of treating infection by HIV.

- 30 Other embodiments, aspects and features of the present invention are either further described in or will be apparent from the ensuing description, examples and appended claims.

DETAILED DESCRIPTION OF THE INVENTION

The present invention includes compounds of Formula I above, and pharmaceutically acceptable salts thereof. These compounds and pharmaceutically acceptable salts thereof are HIV integrase inhibitors.

5 A first embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R^1 is CH_2-R^J ; and all other variables are as originally defined (i.e., as defined in the Summary of the Invention).

A second embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R^1 is CH_2-R^J ; R^J is phenyl, pyridyl, quinolinylnyl, isoquinolinylnyl, cinnolinylnyl, or quinazolinylnyl, any of which is

(a) optionally substituted with from 1 to 4 substituents each of which is independently:

- (1) $-C_{1-4}$ alkyl,
- (2) $-O-C_{1-4}$ alkyl,
- (3) $-C_{1-4}$ haloalkyl,
- 15 (4) $-O-C_{1-4}$ haloalkyl,

- (5) halo,
- (6) $-CN$,
- (7) $-N(RA)RB$,
- (8) $-C(=O)N(RA)RB$,
- 20 (9) $-S(=O)RA$,
- (10) $-SO_2RA$,
- (11) $-N(RA)SO_2RB$,
- (12) $-N(RA)SO_2N(RA)RB$,
- (13) $-N(RA)C(=O)RB$, or
- 25 (14) $-N(RA)C(=O)-C(=O)N(RA)RB$, and

(b) optionally substituted with phenyl, benzyl, $-HetA$, or $-C(=O)-HetA$; wherein each $HetA$ is independently a C_{4-7} azacycloalkyl or a C_{3-6} diazacycloalkyl, either of which is optionally substituted with from 1 to 3 substituents each of which is independently oxo or C_{1-4} alkyl; and with the proviso that when $HetA$ is attached to the rest of the

30 compound via the $-C(=O)-$ moiety, the $HetA$ is attached to the $-C(=O)-$ via a ring N atom; and all other variables are as originally defined.

A third embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R^1 is CH_2-R^J ; R^J is phenyl optionally substituted with from 1 to 3 substituents each of which is independently: (1) $-C_{1-4}$ alkyl, (2) $-C_{1-4}$ fluoroalkyl, (3)

-O-C₁₋₄ alkyl, (4) -O-C₁₋₄ fluoroalkyl, (5) halo, (6) -CN, (7) -C(=O)N(R^A)R^B, or (8) -SO₂R^A; and all other variables are as originally defined.

A fourth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R¹ is CH₂-R^J; R^J is 4-fluorophenyl; and all other

5 variables are as originally defined.

A fifth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R², R³, R⁴ and R⁵ are defined as follows:

(A) R² and R⁴ are as originally defined in part A of the above definition of R², R³, R⁴ and R⁵; and R³ and R⁵ are both H;

10 (B) R² and R⁴ are as originally defined in part B of the above definition of R², R³, R⁴ and R⁵; and R³ and R⁵ are both H;

(C) R² and R³ are as originally defined in part C of the above definition of R², R³, R⁴ and R⁵; and R⁴ and R⁵ are both H; or

15 (D) R⁴ and R⁵ are as originally defined in part D of the above definition of R², R³, R⁴ and R⁵; and R² and R³ are both H;

and all other variables are as originally defined or as defined in any one of the preceding embodiments.

A sixth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R², R³, R⁴ and R⁵ are defined as follows:

20 (A) R³ and R⁵ are both H; and R² and R⁴ are each independently (1) -H, (2) -C₁₋₄ alkyl, (3) -C₁₋₄ fluoroalkyl, (4) C₃₋₆ cycloalkyl, (5) phenyl, or (6) benzyl;

(B) R² and R⁴ together with the carbon atoms to which each is attached form a carbon-carbon double bond; and R³ and R⁵ are both H;

(C) R² and R³ together with the carbon atom to which they are both attached form a 3- to 7-membered saturated carbocyclic ring; and R⁴ and R⁵ are both H; or

25 (D) R⁴ and R⁵ together with the carbon atom to which they are both attached form a 3- to 7-membered saturated carbocyclic ring; and R² and R³ are both H;

and all other variables are as originally defined or as defined in any one of the preceding embodiments.

A seventh embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R², R³, R⁴ and R⁵ are all H; and all other variables are as originally defined or as defined in any one of the preceding embodiments.

30

An eighth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R⁶ is: (1) -H, (2) -C₁₋₆ alkyl, (3) -C₁₋₆ fluoroalkyl, (4) CycA, (5) AryA, or (6) -C₁₋₆ alkyl substituted with AryA; and all other variables are as originally defined or as defined in any one of the preceding embodiments.

A ninth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R^6 is H, $-C_{1-4}$ alkyl, CF_3 , cyclopropyl, phenyl or benzyl; and all other variables are as originally defined or as defined in any one of the preceding embodiments.

5 A tenth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R^6 is H; and all other variables are as originally defined or as defined in any one of the preceding embodiments.

10 An eleventh embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R^7 and R^8 are each independently: (1) -H, (2) $-C_{1-6}$ alkyl, (3) $-CO_2R^A$, (4) $-C(=O)N(R^A)R^B$, (5) $-R^K$, (6) $-C(=O)-R^K$, (7) $-C(=O)N(R^A)-R^K$, or (8) $-C(=O)N(R^A)-C_{1-6}$ alkylene- R^K ; or alternatively R^7 and R^8 together with the carbon atom to which they are both attached form a 3- to 7-membered saturated carbocyclic ring; and all other variables are as originally defined or as defined in any one of the preceding embodiments.

15 A twelfth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R^7 is H or $-C_{1-4}$ alkyl; and R^8 is: (1) -H, (2) $-C_{1-4}$ alkyl, (3) $-CO_2-C_{1-4}$ alkyl, (4) $-C(=O)NH(C_{1-4}$ alkyl), (5) $-C(=O)N(C_{1-4}$ alkyl) $_2$, (6) CycA,

(7) $-C(=O)-HetE$, wherein HetE is a 4- to 7-membered saturated heterocyclic ring containing at least one carbon atom and from 1 to 4 heteroatoms selected from 1 to 4 N atoms, zero or 1 oxygen atoms, and zero or 1 sulfur atoms, wherein the saturated heterocyclic is
20 optionally substituted with from 1 to 3 substituents each of which is independently oxo or C_{1-4} alkyl; and with the proviso that the saturated heterocyclic is attached to the $-C(=O)-$ via a ring N atom, or
(8) $-C(=O)N(R^A)-(CH_2)_{1-2}-HetF$, wherein HetF is a 5- or 6-membered heteroaromatic ring containing from 1 to 4 heteroatoms independently selected from N, O and S, wherein the
25 heteroaromatic ring is optionally substituted with 1 or 2 substituents each of which is independently a C_{1-4} alkyl;

or alternatively R^7 and R^8 together with the carbon atom to which they are both attached form a 3- to 6-membered saturated carbocyclic ring; and all other variables are as originally defined or as defined in any one of the preceding embodiments.

30 A thirteenth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R^9 is: (1) -H, (2) $-C_{1-6}$ alkyl (3) $-C_{1-6}$ fluoroalkyl, (4) CycA, or (5) $-C_{1-6}$ alkyl substituted with CycA, AryA, or HetC; and all other variables are as originally defined or as defined in any one of the preceding embodiments.

A fourteenth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R⁹ is: (1) -H, (2) -C₁₋₄ alkyl, (3) -CH₂CF₃, (4) -C₃₋₆ cycloalkyl, (5) -CH₂-C₃₋₆ cycloalkyl, or (6) -CH₂-phenyl; and all other variables are as originally defined or as defined in any one of the preceding embodiments.

5 A fifteenth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein R⁹ is H, methyl, ethyl, n-propyl, isopropyl, -CH₂CF₃, cyclopropyl, or -CH₂-cyclopropyl.

10 A sixteenth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein each CycA is independently a C₃₋₇ cycloalkyl, which is optionally substituted with from 1 to 4 substituents each of which is halogen, -OH, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, or -O-C₁₋₆ haloalkyl; and all other variables are as originally defined or as defined in any one of the preceding embodiments.

A seventeenth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein each AryA is independently an aryl, which is

15 (a) optionally substituted with from 1 to 5 substituents each of which is independently -C₁₋₆ alkyl, -C₁₋₆ alkylene-OH, -C₁₋₆ alkylene-O-C₁₋₆ alkyl, -C₁₋₆ alkylene-O-C₁₋₆ haloalkyl, -C₁₋₆ alkylene-N(RA)RB, -C₁₋₆ alkylene-C(=O)N(RA)RB, -C₁₋₆ alkylene-C(=O)RA, -C₁₋₆ alkylene-CO₂RA, -C₁₋₆ alkylene-S(O)_nRA, -O-C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ haloalkyl, -OH, halo, -N(RA)RB, -C(=O)N(RA)RB,

20 -C(=O)RA, -CO₂RA, -S(O)_nRA, or -SO₂N(RA)RB, and

(b) optionally substituted with C₃₋₆ cycloalkyl, phenyl, HetD, -CH₂-C₃₋₆ cycloalkyl, benzyl, or -CH₂-HetD;

and all other variables are as originally defined or as defined in any one of the preceding embodiments.

25 An eighteenth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein each HetC is independently:

- (i) a 4- to 7-membered saturated heterocyclic ring containing at least one carbon atom and from 1 to 4 heteroatoms independently selected from N, O and S, wherein the saturated ring is:
- 30 (a) optionally substituted with from 1 to 4 substituents each of which is halogen, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, OH, or oxo, and
- (b) optionally substituted with C₃₋₆ cycloalkyl, phenyl, HetD, -CH₂-C₃₋₆ cycloalkyl, benzyl, or -CH₂-HetD;
- (ii) a 5- or 6-membered heteroaromatic ring containing from 1 to 4 heteroatoms independently selected from N, O and S, wherein the heteroaromatic ring is:

- (a) optionally substituted with from 1 to 4 substituents each of which is halogen, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, or OH, and
- (b) optionally substituted with C₃₋₆ cycloalkyl, phenyl, HetD, -CH₂-C₃₋₆ cycloalkyl, benzyl, or -CH₂-HetD;

5 and all other variables are as originally defined or as defined in any one of the preceding embodiments.


A nineteenth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein each HetD is independently (i) a 4- to 7-membered saturated heterocyclic ring containing at least one carbon atom and from 1 to 4 heteroatoms independently selected from N, O and S or (ii) a 5- or 6-membered heteroaromatic ring containing from 1 to 4 heteroatoms independently selected from N, O and S; and all other variables are as originally defined or as defined in any one of the preceding embodiments.

A twentieth embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein each aryl is independently phenyl, indenyl, indanyl, naphthyl, or tetrahydronaphthyl; and all other variables are as originally defined or as defined in any one of the preceding embodiments. In an aspect of this embodiment, each aryl is independently phenyl or naphthyl. In another aspect each aryl is phenyl.

A twenty-first embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein each R^A and R^B is independently H or C₁₋₄ alkyl; and all other variables are as originally defined or as defined in any one of the preceding embodiments.

A twenty-second embodiment of the present invention is a compound of Formula I, or a pharmaceutically acceptable salt thereof, wherein each R^A and R^B is independently H, methyl, or ethyl; and all other variables are as originally defined or as defined in any one of the preceding embodiments. In an aspect of this embodiment, each R^A and R^B is independently H or methyl.

A first class of the present invention includes compounds of Formula II, and pharmaceutically acceptable salts thereof:



wherein:

X¹ and X² are each independently -H, -C₁₋₄ alkyl, -O-C₁₋₄ alkyl, -C₁₋₄ haloalkyl, -O-C₁₋₄ haloalkyl, halo, -CN, -N(R^A)R^B, -C(=O)N(R^A)R^B, or -S(O)_nR^A, wherein n is an integer equal to zero, 1, or 2;

R², R³, R⁴ and R⁵ are defined as follows:

- 5 (A) R² and R⁴ are each independently -H, -C₁₋₄ alkyl, -C₁₋₄ fluoroalkyl, C₃₋₆ cycloalkyl, phenyl, or benzyl; and R⁴ and R⁵ are both H;
- (B) R² and R⁴ together with the carbon atoms to which each is attached form a carbon-carbon double bond; and R³ and R⁵ are both H;
- (C) R² and R³ together with the carbon atom to which they are both attached form
10 cyclopropyl; and R⁴ and R⁵ are both H; or
- (D) R⁴ and R⁵ together with the carbon atom to which they are both attached form cyclopropyl; and R² and R³ are both H;

R⁶ is H, -C₁₋₄ alkyl, CF₃, cyclopropyl, phenyl or benzyl;

15

R⁷ is H or -C₁₋₄ alkyl;

R⁸ is -H, -C₁₋₄ alkyl, -CO₂-C₁₋₄ alkyl, -C(=O)NH(C₁₋₄ alkyl), -C(=O)N(C₁₋₄ alkyl)₂, C₃₋₆ cycloalkyl, -C(=O)-HetE, or -C(=O)N(R^A)-(CH₂)₁₋₂-HetF; wherein

20 HetE is a 4- to 7-membered saturated heterocyclic ring containing at least one carbon atom and from 1 to 4 heteroatoms selected from 1 to 4 N atoms, zero or 1 oxygen atoms, and zero or 1 sulfur atoms, wherein the saturated heterocyclic is optionally substituted with from 1 to 3 substituents each of which is independently oxo or C₁₋₄ alkyl; and with the proviso that the saturated heterocyclic is attached to the -C(=O)- via a ring N atom; and

25 HetF is a 5- or 6-membered heteroaromatic ring containing from 1 to 4 heteroatoms independently selected from N, O and S, wherein the heteroaromatic ring is optionally substituted with 1 or 2 substituents each of which is independently a C₁₋₄ alkyl;

or alternatively R⁷ and R⁸ together with the carbon atom to which they are both attached form a 3- to 6-
30 membered saturated carbocyclic ring;

R⁹ is -H, -C₁₋₄ alkyl, -CH₂CF₃, -C₃₋₆ cycloalkyl, -CH₂-C₃₋₆ cycloalkyl, or -CH₂-phenyl;

each R^A is independently H or C₁₋₄ alkyl; and

each R^B is independently H or C₁₋₄ alkyl.

5 A sub-class of the first class includes compounds of Formula II, and pharmaceutically acceptable salts thereof, wherein:

X¹ and X² are each independently H, fluoro, chloro, methyl, trifluoromethyl, methoxy, CN, -SO₂CH₃, -C(=O)NH(CH₃), or -C(=O)N(CH₃)₂;


10 R², R³, R⁴ and R⁵ are all H;



R⁶ is H, methyl, cyclopropyl, or phenyl;

R⁷ is H or methyl;

15 R⁸ is -H, -C₁₋₄ alkyl, -CO₂-C₁₋₄ alkyl, -C(=O)NH(C₁₋₄ alkyl), -C(=O)N(C₁₋₄ alkyl)₂, C₃₋₆ cycloalkyl, -C(=O)-HetE, or -C(=O)N(R^A)-(CH₂)₁₋₂-HetF; wherein

HetE is selected from the group consisting of:



, and , wherein the asterisk * denotes the point of attachment to the -C(=O) moiety; and

HetF is selected from the group consisting of pyrrolyl, imidazolyl, triazolyl, tetrazolyl, oxazolyl, isooxazolyl, pyridyl, pyrimidinyl, and pyrazinyl;

25 or alternatively R⁷ and R⁸ together with the carbon atom to which they are both attached form cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; and

R⁹ is H, methyl, ethyl, n-propyl, isopropyl, -CH₂CF₃, cyclopropyl, or -CH₂-cyclopropyl.

30 Another embodiment of the present invention is a compound, or a pharmaceutically acceptable salt thereof, selected from the group consisting of the compounds set forth in Examples 1 to 16 below.

Other embodiments of the present invention include the following:

(a) A pharmaceutical composition comprising an effective amount of a compound of Formula I and a pharmaceutically acceptable carrier.

5 (b) A pharmaceutical composition which comprises the product prepared by combining (e.g., mixing) an effective amount of a compound of Formula I and a pharmaceutically acceptable carrier.

(c) The pharmaceutical composition of (a) or (b), further comprising an effective amount of an HIV infection/AIDS treatment agent selected from the group consisting of HIV/AIDS antiviral agents, immunomodulators, and anti-infective agents.

10 (d) The pharmaceutical composition of (c), wherein the HIV infection/AIDS treatment agent is an antiviral selected from the group consisting of HIV protease inhibitors, non-nucleoside HIV reverse transcriptase inhibitors, and nucleoside HIV reverse transcriptase inhibitors.

(e) A pharmaceutical combination which is (i) a compound of Formula I and (ii) an HIV infection/AIDS treatment agent selected from the group consisting of HIV/AIDS antiviral agents, immunomodulators, and anti-infective agents; wherein the compound of Formula I and the HIV infection/AIDS treatment agent are each employed in an amount that renders the combination effective for inhibiting HIV integrase, for treating or preventing infection by HIV, or for preventing, treating or delaying the onset of AIDS.

20 (f) The combination of (e), wherein the HIV infection/AIDS treatment agent is an antiviral selected from the group consisting of HIV protease inhibitors, non-nucleoside HIV reverse transcriptase inhibitors and nucleoside HIV reverse transcriptase inhibitors.

(g) A method of inhibiting HIV integrase in a subject in need thereof which comprises administering to the subject an effective amount of a compound of Formula I.

25 (h) A method of preventing or treating infection by HIV in a subject in need thereof which comprises administering to the subject an effective amount of a compound of Formula I.

(i) The method of (h), wherein the compound of Formula (I) is administered in combination with an effective amount of at least one antiviral selected from the group consisting of HIV protease inhibitors, non-nucleoside HIV reverse transcriptase inhibitors, and nucleoside HIV reverse transcriptase inhibitors.

30 (j) A method of preventing, treating or delaying the onset of AIDS in a subject in need thereof which comprises administering to the subject an effective amount of a compound of Formula I.

(k) The method of (j), wherein the compound is administered in combination with an effective amount of at least one antiviral selected from the group consisting of HIV protease

inhibitors, non-nucleoside HIV reverse transcriptase inhibitors, and nucleoside HIV reverse transcriptase inhibitors

(l) A method of inhibiting HIV integrase in a subject in need thereof which comprises administering to the subject the pharmaceutical composition of (a), (b), (c) or (d) or the combination of (e) or (f).

(m) A method of preventing or treating infection by HIV in a subject in need thereof which comprises administering to the subject the pharmaceutical composition of (a), (b), (c) or (d) or the combination of (e) or (f).

(n) A method of preventing, treating or delaying the onset of AIDS in a subject in need thereof which comprises administering to the subject the pharmaceutical composition of (a), (b), (c) or (d) or the combination of (e) or (f).

The present invention also includes a compound of the present invention (i) for use in, (ii) for use as a medicament for, or (iii) for use in the preparation of a medicament for: (a) inhibiting HIV integrase, (b) preventing or treating infection by HIV, or (c) preventing, treating or delaying the onset of AIDS. In these uses, the compounds of the present invention can optionally be employed in combination with one or more HIV/AIDS treatment agents selected from HIV/AIDS antiviral agents, anti-infective agents, and immunomodulators.

Additional embodiments of the invention include the pharmaceutical compositions, combinations and methods set forth in (a)-(n) above and the uses set forth in the preceding paragraph, wherein the compound of the present invention employed therein is a compound of one of the embodiments, aspects, classes, sub-classes, or features of the compounds described above. In all of these embodiments, the compound may optionally be used in the form of a pharmaceutically acceptable salt.

As used herein, the term "alkyl" refers to any linear or branched chain alkyl group having a number of carbon atoms in the specified range. Thus, for example, "C₁₋₆ alkyl" (or "C₁-C₆ alkyl") refers to all of the hexyl alkyl and pentyl alkyl isomers as well as n-, iso-, sec- and t-butyl, n- and isopropyl, ethyl and methyl. As another example, "C₁₋₄ alkyl" refers to n-, iso-, sec- and t-butyl, n- and isopropyl, ethyl and methyl.

The term "alkylene" refers to any linear or branched chain alkylene group (or alternatively "alkanediyl") having a number of carbon atoms in the specified range. Thus, for example, "-C₁₋₆ alkylene-" refers to any of the C₁ to C₆ linear or branched alkylenes. A class of alkylenes of particular interest with respect to the invention is -(CH₂)₁₋₆-, and sub-classes of particular interest include -(CH₂)₁₋₄-, -(CH₂)₁₋₃-, -(CH₂)₁₋₂-, and -CH₂-. Also of interest is the alkylene -CH(CH₃)-

The term "cycloalkyl" refers to any cyclic ring of an alkane having a number of carbon atoms in the specified range. Thus, for example, "C₃₋₈ cycloalkyl" (or "C₃-C₈ cycloalkyl") refers to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

5 The term "halogen" (or "halo") refers to fluorine, chlorine, bromine and iodine (alternatively referred to as fluoro, chloro, bromo, and iodo).

The term "haloalkyl" refers to an alkyl group as defined above in which one or more of the hydrogen atoms has been replaced with a halogen (i.e., F, Cl, Br and/or I). Thus, for example, "C₁₋₆ haloalkyl" (or "C₁-C₆ haloalkyl") refers to a C₁ to C₆ linear or branched alkyl group as defined above with one or more halogen substituents. The term "fluoroalkyl" has an analogous meaning except that the
10 halogen substituents are restricted to fluoro. Suitable fluoroalkyls include the series (CH₂)₀₋₄CF₃ (i.e., trifluoromethyl, 2,2,2-trifluoroethyl, 3,3,3-trifluoro-n-propyl, etc.).

The term "C₄₋₇ azacycloalkyl" (or "C₄-C₇ azacycloalkyl") means a saturated cyclic ring consisting of one nitrogen and from four to seven carbon atoms (i.e., pyrrolidinyl, piperidinyl, azepanyl, or octahydroazocinyl).
15

The term "C₃₋₆ diazacycloalkyl" (or "C₃-C₆ diazacycloalkyl") means a saturated cyclic ring consisting of two nitrogens and from three to six carbon atoms (e.g., imidazolidinyl, pyrazolidinyl, or piperazinyl).
20

Unless expressly stated to the contrary, all ranges cited herein are inclusive. For example, a heterocyclic ring described as containing from "1 to 4 heteroatoms" means the ring can
20 contain 1, 2, 3 or 4 heteroatoms. It is also to be understood that any range cited herein includes within its scope all of the sub-ranges within that range. Thus, for example, a heterocyclic ring described as containing from "1 to 4 heteroatoms" is intended to include as aspects thereof, heterocyclic rings containing 2 to 4 heteroatoms, 3 or 4 heteroatoms, 1 to 3 heteroatoms, 2 or 3 heteroatoms, 1 or 2 heteroatoms, 1 heteroatom, 2 heteroatoms, and so forth.


25 When any variable (e.g., R^A and R^B) occurs more than one time in any constituent or in Formula I, Formula II, or in any other formula depicting and describing compounds of the invention, its definition on each occurrence is independent of its definition at every other occurrence. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

30 The term "substituted" (e.g., as in "is optionally substituted with from 1 to 5 substituents ...") includes mono- and poly-substitution by a named substituent to the extent such single and multiple substitution (including multiple substitution at the same site) is chemically allowed. Unless expressly stated to the contrary, substitution by a named substituent is permitted on any atom in a ring (e.g.,

cycloalkyl, aryl, a heteroaromatic ring, or a saturated heterocyclic ring) provided such ring substitution is chemically allowed and results in a stable compound.

Unless expressly stated to the contrary, an “unsaturated” ring is a partially or fully unsaturated ring.

- 5 In instances where a hydroxy (-OH) substituent(s) is(are) permitted on a heteroaromatic ring and keto-enol tautomerism is possible, it is understood that the substituent might in fact be present, in whole or in part, in the keto form, as exemplified here for a hydroxypyridinyl substituent:




- 10 Compounds of the present invention having a hydroxy substituent on a carbon atom of a heteroaromatic ring are understood to include compounds in which only the hydroxy is present, compounds in which only the tautomeric keto form (i.e., an oxo substituent) is present, and compounds in which the keto and enol forms are both present. Similarly, where a hydroxy substituent is permitted on an unsaturated, non-aromatic heterocyclic ring and keto-enol tautomerism is possible, or an oxo substituent is permitted on a saturated heterocyclic ring and keto-enol tautomerism is possible, it is understood that the substituent
- 15 might in fact be present as the keto tautomer, the enol tautomer or a mixture thereof. It is understood that compounds of the invention described (e.g., in Formula I) in terms of either the keto form or the enol form include compounds in which either or both the keto and enol forms are present.

- Any of the various carbocyclic and heterocyclic rings and ring systems defined herein may be attached to the rest of the compound at any ring atom (i.e., any carbon atom or any heteroatom)
- 20 provided that a stable compound results. A class of cycloalkyl groups suitable for use in the compounds of the invention (e.g., in the definition of CycA) consists of the C₃₋₆ cycloalkyl groups - cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. A class of aryl groups suitable for use in the invention (e.g., independently in the definitions of either or both R^J and AryA) consists of phenyl, indenyl, indanyl, naphthyl, and tetrahydronaphthyl. A sub-class of aryl groups particularly suitable for use in the present
- 25 invention consists of phenyl and naphthyl. Another aryl sub-class of particular interest is phenyl. A class of 4- to 7-membered saturated heterocyclic rings suitable for use in the present invention (e.g., independently in the definitions of one or more of HetC, HetD and HetE) consists of piperidinyl, morpholinyl, thiomorpholinyl, thiazolidinyl, isothiazolidinyl, oxazolidinyl, isoxazolidinyl, pyrrolidinyl, azetidiny, imidazolidinyl, piperazinyl, tetrahydrofuranyl, tetrahydrothienyl, pyrazolidinyl,

hexahydropyrimidinyl, thiazinanyl, thiazepanyl, thiadiazepanyl, dithiazepanyl, azepanyl, diazepanyl, thiadiazinanyl, tetrahydropyranyl, tetrahydrothiopyranyl, and dioxanyl. A class of 4- to 7-membered unsaturated heterocyclic rings suitable for use in the present invention (e.g., independently in the definitions of either or both HetC and HetD) consists of the mono-unsaturated counterparts of the class of saturated heterocyclic rings set forth in the preceding sentence. A class of 5- or 6-membered heteroaromatic rings suitable for use in the present invention (e.g., independently in any one or more of the definitions of R^J, HetB, HetC, HetD and HetF) consists of pyridyl, pyrrolyl, pyrazinyl, pyrimidinyl, pyridazinyl, triazinyl, thienyl, furanyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, oxazolyl, isooxazolyl, oxadiazolyl, oxatriazolyl, thiazolyl, isothiazolyl, and thiadiazolyl.

A "stable" compound is a compound which can be prepared and isolated and whose structure and properties remain or can be caused to remain essentially unchanged for a period of time sufficient to allow use of the compound for the purposes described herein (e.g., therapeutic or prophylactic administration to a subject).


As would be recognized by one of ordinary skill in the art, the compounds of the present invention can exist as tautomers such as the following:



For the purposes of the present invention a reference herein to a compound of Formula I or Formula II is a reference to compound I or compound II per se, or to any one of its tautomers per se, or to mixtures thereof.

The compounds of the present invention have at least one asymmetric center at the fused

ring carbon in the naphthyridine ring marked by the arrow in Formula I:



Additional asymmetric centers may be present depending upon the nature of other substituents in the molecule. Each such asymmetric center will independently produce two optical isomers. All possible optical isomers and diastereomers of these compounds, individually and in mixtures, are within the scope of the present invention.

The compounds of the present inventions are useful in the inhibition of HIV integrase, the prevention or treatment of infection by human immunodeficiency virus (HIV) and the prevention, treatment or the delay in the onset of consequent pathological conditions such as AIDS. Preventing AIDS, treating AIDS, delaying the onset of AIDS, or preventing or treating infection by HIV is defined as including, but not limited to, treatment of a wide range of states of HIV infection: AIDS, ARC (AIDS related complex), both symptomatic and asymptomatic, and actual or potential exposure to HIV. For example, the compounds of this invention are useful in treating infection by HIV after suspected past exposure to HIV by such means as blood transfusion, exchange of body fluids, bites, accidental needle stick, or exposure to patient blood during surgery.

The compounds of this invention are useful in the preparation and execution of screening assays for antiviral compounds. For example, the compounds of this invention are useful for isolating enzyme mutants, which are excellent screening tools for more powerful antiviral compounds. Furthermore, the compounds of this invention are useful in establishing or determining the binding site of other antivirals to HIV integrase, e.g., by competitive inhibition. Thus the compounds of this invention are commercial products to be sold for these purposes.

The compounds of the present invention may be administered in the form of pharmaceutically acceptable salts. The term "pharmaceutically acceptable salt" refers to a salt which possesses the effectiveness of the parent compound and which is not biologically or otherwise undesirable (e.g., is neither toxic nor otherwise deleterious to the recipient thereof). Suitable salts include acid addition salts which may, for example, be formed by mixing a solution of the compound of the present invention with a solution of a pharmaceutically acceptable acid such as hydrochloric acid, sulfuric acid, acetic acid, trifluoroacetic acid, or benzoic acid. Many of the compounds of the invention carry an acidic moiety, in which case suitable pharmaceutically acceptable salts thereof can include alkali

metal salts (e.g., sodium or potassium salts), alkaline earth metal salts (e.g., calcium or magnesium salts), and salts formed with suitable organic ligands such as quaternary ammonium salts. Also, in the case of an acid (-COOH) or alcohol group being present, pharmaceutically acceptable esters can be employed to modify the solubility or hydrolysis characteristics of the compound.

5 The term "administration" and variants thereof (e.g., "administering" a compound) in reference to a compound of the invention mean providing the compound or a prodrug of the compound to the individual in need of treatment. When a compound of the invention or a prodrug thereof is provided in combination with one or more other active agents (e.g., antiviral agents useful for treating HIV infection or AIDS), "administration" and its variants are each understood to include concurrent and
10 sequential provision of the compound or prodrug and other agents.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combining the specified ingredients in the specified amounts.

By "pharmaceutically acceptable" is meant that the ingredients of the pharmaceutical
15 composition must be compatible with each other and not deleterious to the recipient thereof.

The term "subject" (alternatively referred to herein as "patient") as used herein refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment.

The term "effective amount" as used herein means that amount of active compound or
20 pharmaceutical agent that elicits the biological or medicinal response in a tissue, system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician. In one embodiment, the effective amount is a "therapeutically effective amount" for the alleviation of the symptoms of the disease or condition being treated. In another embodiment, the effective amount is a "prophylactically effective amount" for prophylaxis of the symptoms of the disease or condition being
25 prevented. The term also includes herein the amount of active compound sufficient to inhibit HIV integrase and thereby elicit the response being sought (i.e., an "inhibition effective amount"). When the active compound (i.e., active ingredient) is administered as the salt, references to the amount of active ingredient are to the free acid or free base form of the compound.

For the purpose of inhibiting HIV integrase, preventing or treating HIV infection or
30 preventing, treating or delaying the onset of AIDS, the compounds of the present invention, optionally in the form of a salt, can be administered by any means that produces contact of the active agent with the agent's site of action. They can be administered by any conventional means available for use in conjunction with pharmaceuticals, either as individual therapeutic agents or in a combination of therapeutic agents. They can be administered alone, but typically are administered with a pharmaceutical

carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice. The compounds of the invention can, for example, be administered orally, parenterally (including subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques), by inhalation spray, or rectally, in the form of a unit dosage of a pharmaceutical composition containing an effective amount of the compound and conventional non-toxic pharmaceutically-acceptable carriers, adjuvants and vehicles. Liquid preparations suitable for oral administration (e.g., suspensions, syrups, elixirs and the like) can be prepared according to techniques known in the art and can employ any of the usual media such as water, glycols, oils, alcohols and the like. Solid preparations suitable for oral administration (e.g., powders, pills, capsules and tablets) can be prepared according to techniques known in the art and can employ such solid excipients as starches, sugars, kaolin, lubricants, binders, disintegrating agents and the like. Parenteral compositions can be prepared according to techniques known in the art and typically employ sterile water as a carrier and optionally other ingredients, such as a solubility aid. Injectable solutions can be prepared according to methods known in the art wherein the carrier comprises a saline solution, a glucose solution or a solution containing a mixture of saline and glucose. Further description of methods suitable for use in preparing pharmaceutical compositions of the present invention and of ingredients suitable for use in said compositions is provided in Remington's Pharmaceutical Sciences, 18th edition, edited by A. R. Gennaro, Mack Publishing Co., 1990.

The compounds of this invention can be administered orally in a dosage range of 0.001 to 1000 mg/kg of mammal (e.g., human) body weight per day in a single dose or in divided doses. One preferred dosage range is 0.01 to 500 mg/kg body weight per day orally in a single dose or in divided doses. Another preferred dosage range is 0.1 to 100 mg/kg body weight per day orally in single or divided doses. For oral administration, the compositions can be provided in the form of tablets or capsules containing 1.0 to 500 milligrams of the active ingredient, particularly 1, 5, 10, 15, 20, 25, 50, 75, 100, 150, 200, 250, 300, 400, and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. The specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, general health, sex, diet, mode and time of administration, rate of excretion, drug combination, the severity of the particular condition, and the host undergoing therapy.

As noted above, the present invention is also directed to use of the HIV integrase inhibitor compounds of the present invention with one or more agents useful in the treatment of HIV infection or AIDS. For example, the compounds of this invention may be effectively administered, whether at periods of pre-exposure and/or post-exposure, in combination with effective amounts of one or more HIV/AIDS antivirals, immunomodulators, antiinfectives, or vaccines useful for treating HIV

infection or AIDS, such as those disclosed in Table 1 of WO 01/38332 or in the Table in WO 02/30930. Suitable HIV/AIDS antivirals for use in combination with the compounds of the present invention include, for example, HIV protease inhibitors (e.g., indinavir, atazanavir, lopinavir optionally with ritonavir, saquinavir, or nelfinavir), nucleoside HIV reverse transcriptase inhibitors (e.g., abacavir, lamivudine (3TC), zidovudine (AZT), or tenofovir), and non-nucleoside HIV reverse transcriptase inhibitors (e.g., efavirenz or nevirapine). It will be understood that the scope of combinations of the compounds of this invention with HIV/AIDS antivirals, immunomodulators, anti-infectives or vaccines is not limited to the foregoing substances or to the list in the above-referenced Tables in WO 01/38332 and WO 02/30930, but includes in principle any combination with any pharmaceutical composition useful for the treatment of AIDS. The HIV/AIDS antivirals and other agents will typically be employed in these combinations in their conventional dosage ranges and regimens as reported in the art, including, for example, the dosages described in the Physicians' Desk Reference, 57th edition, Thomson PDR, 2003. The dosage ranges for a compound of the invention in these combinations are the same as those set forth above.

Abbreviations used in the instant specification, particularly the in the Schemes and Examples, include the following:

AIDS = acquired immunodeficiency syndrome
 ARC = AIDS related complex
 DBU = 1,8-diazabicyclo[5.4.0]undec-7-ene
 DIEA = diisopropylethylamine (or Hunig's base)
 DMF = N,N-dimethylformamide
 DMSO = dimethylsulfoxide
 EDC = 1-ethyl-3-(3-dimethylaminopropyl) carbodiimide
 ES MS = electrospray mass spectroscopy
 Et = ethyl
 EtOAc = ethyl acetate
 HIV = human immunodeficiency virus
 HOBT or HOBt = 1-hydroxy benzotriazole hydrate
 HPLC = high performance liquid chromatography
 i-Pr = isopropyl
 LiHMDS = lithium hexamethyldisilazide
 Me = methyl
 NMR = nuclear magnetic resonance
 TFA = trifluoroacetic acid


THF = tetrahydrofuran

The compounds of the present invention can be readily prepared according to the following reaction schemes and examples, or modifications thereof, using readily available starting materials, reagents and conventional synthesis procedures. In these reactions, it is also possible to make use of variants which are themselves known to those of ordinary skill in this art, but are not mentioned in greater detail. Furthermore, other methods for preparing compounds of the invention will be readily apparent to the person of ordinary skill in the art in light of the following reaction schemes and examples. Unless otherwise indicated, all variables are as defined above.

Schemes 1 to 3 present methods for preparing compounds of the present invention which contain the 2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione bicyclic nucleus. In Scheme 1, 2-piperidinone 1 is N-alkylated with a halide compound or equivalent reagent in the presence of a base such as sodium hydride to give derivative 2 which contains the R¹ variable. Sulfinylation of 2 using methyl phenylsulfinate in the presence of a strong base such as lithium hexamethyldisilazide, followed by heating the sulfinate product in the presence of a base such as sodium carbonate gives unsaturated lactam 3. Michael addition of a nitro compound to 3 in the presence of a base such as 1,8-diazabicyclo[5.4.0]undec-7-ene gives the 4-substituted piperidinone 4, the nitro group in which is reduced to amine 5 using a reagent such as Raney nickel in the presence of hydrogen gas. There are then several pathways to convert 5 to the desired 2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione products. One method involves acylation of 5 with methyl oxalyl chloride or an equivalent reagent to give 6. Oxalamide 6 is then cyclized to 2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione 7 in the presence of a strong base such as lithium hexamethyldisilazide. Introducing the R⁹ variable on the nitrogen at position 6 in compound 7 is accomplished in an alkylation reaction using a halide compound or an equivalent reagent and a base such as cesium carbonate. During this reaction, in addition to the desired alkylation of the nitrogen at position 6, the hydroxyl group at position 8 may also become alkylated. Removal of the alkyl group on oxygen is achieved in a second step using a reagent such as hydrogen bromide to give compound 10. In a second method, oxalamide 6 is converted to 2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione 10 in a one-pot procedure which involves treatment of 6 with a strong base such as lithium hexamethyldisilazide and a halide compound or equivalent reagent to install the R⁹ variable, followed by the addition of more strong base such as lithium hexamethyldisilazide to close the ring and furnish 10. In a third method, the amino group in 5 is alkylated with a halide compound or an equivalent reagent to introduce the R⁹ variable, giving 8. Amine 8 is treated with two or more equivalents of a strong base such as lithium diisopropylamide, and the resulting anion is treated with diethyl oxalate or equivalent reagent to provide 2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione 10. In a fourth method, compound 8 is acylated with methyl oxalyl chloride or equivalent reagent to give

9. Oxalamide **9** is then cyclized to 2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione **10** in the presence of a strong base such as lithium hexamethyldisilazide.

Scheme 1




In Scheme 2, nitrile **11** is alkylated with allyl bromide or an equivalent reagent in the presence of a strong base such as lithium diisopropylamide. The product from this reaction is then alkylated with methyl bromoacetate or an equivalent reagent in the presence of a strong base such as lithium diisopropylamide to give olefin ester **12**. The ester group in **12** is then converted to an amide by

first hydrolyzing the ester to an acid with a base such as sodium hydroxide in water, then the acid is coupled to an amino compound containing the R¹ variable using a reagent such as EDC to give olefin amide 13. The olefin in 13 is then oxidatively cleaved using a reagent such as ozone to give an aldehyde which cyclizes onto the amide nitrogen with dehydration under acid catalysis to give dihydropyridinone

5 14. The double bond in 14 is reduced with a reagent such as hydrogen in the presence of palladium on carbon to give 15. The nitrile in 15 is converted to thioamide 16 using hydrogen sulfide in the presence of a base such as pyridine. Thioamide 16 is then reduced to amine 17 using a reagent such as Raney nickel activated with sodium hydroxide. Amine 17 is treated with methyl oxalyl chloride or an equivalent reagent to give 18. Oxalamide 18 is then treated with a strong base such as lithium


10 hexamethyldisilazide and a halide compound or an equivalent reagent to install the R⁹ variable, followed by the addition of more strong base such as lithium hexamethyldisilazide to close the ring and furnish 2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione 19.

Scheme 2



- In Scheme 3, an amino compound containing the R^9 variable is alkylated with tert-butyl bromoacetate **20** or an equivalent reagent to give **21**. Glycine ester **21** is then acylated with methyl oxalyl chloride or an equivalent reagent to provide **22**. Oxalamide **22** is combined with unsaturated lactam **3** in the presence of a strong base such as lithium hexamethyldisilazide which induces a tandem Michael addition and Dieckman-type ring closure to give 2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione **23**. The ester group in **23** can then be converted to 2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione amide derivatives **24** in a two-step process involving conversion of the tert-butyl ester to the carboxylic acid using an acid such as trifluoroacetic acid, followed by activation and coupling of the carboxylic acid to an amino compound.


Scheme 3



Schemes 4 and 5 show methods to make 2-piperidinones which incorporate R² and R⁴ substituents on the piperidinone ring. The 2-piperidinone products **28** and **32** in Schemes 4 and 5 can be used as starting materials in place of compound **2** in Scheme 1 to provide 2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione products bearing R² and R⁴ substituents at positions 3 and 4 on the 2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione ring.


In Scheme 4, delta-keto ester derivative **25** containing the R² variable is treated with hydroxylamine to give oxime **26**. The oxime group in **26** is converted to an amino group under reducing conditions such as palladium on carbon and hydrogen gas. The resulting amino group then cyclizes onto the ester to form piperidinone **27**. Compound **27** is then N-alkylated with a halide compound or an equivalent reagent in the presence of a base such as sodium hydride to give piperidinone **28** which contains the R¹ variable and the R² variable at position 6 on the piperidinone ring. Alternatively, the keto group in delta-keto ester derivative **25** is reductively aminated with an amino compound containing the R¹ variable in the presence of a reducing agent such as sodium cyanoborohydride. The resulting amine then cyclizes onto the ester group to give the substituted piperidinone **28**.

Scheme 4




In Scheme 5, nitrile **29** containing the R^4 variable is deprotonated with a strong base such as lithium diisopropylamide and the resulting anion undergoes Michael addition to ethyl acrylate or an equivalent reagent to provide **30**. The nitrile group in **30** is converted to an amine using reductive conditions such as palladium on carbon and hydrogen gas, and the amine then cyclizes onto the ester to form piperidinone **31**. Compound **31** is then N-alkylated with a halide compound or an equivalent reagent in the presence of a base such as sodium hydride to give piperidinone **32** which contains the R^1 variable and the R^4 variable at position 5 on the piperidinone ring.

Scheme 5




Schemes 6 and 7 present methods for preparing compounds of the present invention which contain 2,4a,5,6-tetrahydro-2,6-naphthyridine-1,7-dione bicyclic nucleus. In Scheme 6, the nitrile group in dihydropyridinone **14** from Scheme 2 is reacted with an organometallic reagent such as an organocerium reagent to introduce R^7 and R^8 variables and give **33**. Amine **33** is acylated with methyl oxalyl chloride or equivalent reagent to give **34**. Oxalamide **34** is then treated with a strong base such as lithium hexamethyldisilazide and a halide compound or an equivalent reagent to install the R^9 variable, followed by the addition of more strong base such as lithium hexamethyldisilazide to close the ring and furnish 2,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione **35**.

Scheme 6



In Scheme 7, enediester **36** undergoes Michael addition of a nitro compound containing the R^7 and R^8 variables in the presence of a base such as DBU to give **37**. The two ester groups in **37** are hydrolyzed to give a diacid, and the diacid is treated with a reagent such as acetic anhydride to close the ring to give **38**. Anhydride **38** is then used to acylate an amino compound containing the R^1 variable to give an amide acid, treatment of which with a reagent such as EDC closes the ring to give **39**. Imide **39** is treated with a reducing agent such as sodium borohydride to form a hydroxy piperidinone which undergoes dehydration upon workup in the presence of an acid such as hydrogen chloride to give dihydropyridinone **40**. The nitro group in **40** is reduced with a reagent such as Raney nickel activated with sodium hydroxide to give **41**. Amine **41** is acylated with methyl oxalyl chloride or equivalent reagent to give **42**. Oxalamide **42** is then treated with a strong base such as lithium hexamethyldisilazide and a halide compound or an equivalent reagent to install the R^9 variable, followed by the addition of more strong base such as lithium hexamethyldisilazide to close the ring and provide 2,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione **43**.


Scheme 7



5 The following examples serve only to illustrate the invention and its practice. The examples are not to be construed as limitations on the scope or spirit of the invention.

EXAMPLE 1

2-(4-Fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione



10 Step 1: 1-(4-Fluorobenzyl)piperidine-2-one.

To a solution of 4-fluorobenzyl bromide (15 g, 79 mmol) and 2-piperidinone (8.4 g, 85 mmol) in DMF (150 mL) was added sodium hydride (3.6 g of a 60% suspension in mineral oil, 90 mmol) in portions over a period of 20 min. The mixture was stirred at ambient temperature for 18 hours. The solvent was removed under vacuum and the residue was partitioned between EtOAc and water. The organic extract was washed with brine, dried over MgSO₄, filtered, and concentrated under vacuum. The residual oil was purified using silica gel column chromatography eluting with 50 % - 70 % ethyl acetate in hexanes. The appropriate fractions were combined and concentrated to afford the title compound as white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.23 (dd, *J* = 8.7; 5.4 Hz, 2H), 7.00 (t, *J* = 8.7 Hz, 2H), 4.56 (s, 2H), 3.18 (t, *J* = 6 Hz, 2H), 2.46 (t, *J* = 6 Hz, 2H), 1.79 (m, 4H).

Step 2: 1-(4-Fluorobenzyl)-3-(phenylsulfinyl)piperidin-2-one.

To a cooled (-20 °C) solution of 1-(4-fluorobenzyl)piperidine (17 g, 82 mmol) in anhydrous THF (200 mL) was added LiHMDS (1.0M in THF, 180 mL, 180 mmol) over a period of 15 min. The reaction mixture was stirred for 30 min at -20 °C, then methyl phenylsulfinate (15 g, 98 mmol) was added over a period of 5 min. After being stirred at -20 °C for 1 hour, the reaction was quenched by the addition of water (100 mL). EtOAc (200 mL) was added and the organic layer was separated and washed with water (3 x 100 mL) and brine. The organic layer was dried over Mg₂SO₄, filtered, concentrated under vacuum. The crude product was used without purification in the next step.

Step 3: 1-(4-Fluorobenzyl)-5,6-dihydropyridin-2-(1*H*)-one.

To a stirred solution of 1-(4-fluorobenzyl)-3-(phenylsulfinyl)piperidin-2-one (25 g, 74 mmol) in toluene (200 mL) was added anhydrous sodium carbonate (20 g, 190 mmol). The mixture was heated to reflux for 1.5 hours. The reaction mixture was cooled to ambient temperature and the solids were removed by filtration. The filtrate solvent was removed under vacuum and the residue was purified using silica gel column chromatography eluting with 3:1 EtOAc:hexanes. Appropriate fractions were combined and removal of the solvents under vacuum gave the title compound as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.26 (m, 2H), 7.01 (m, 2H), 6.56 (dt, *J* = 9.9, 4.2 Hz, 1H), 6.00 (dt, *J* = 9.7, 1.8 Hz, 1H), 4.59 (s, 2H), 3.32 (t, *J* = 7.2 Hz, 2H), 2.33 (m, 2H).

Step 4: 1-(4-Fluorobenzyl)-4-(nitromethyl)piperidin-2-one.

To a solution of 1-(4-fluorobenzyl)-5,6-dihydropyridin-2-(1*H*)-one (6.2 g, 30 mmol) in nitromethane (64.5 g, 1.06 mol) under an atmosphere of nitrogen was added DBU (4.59 g, 30.2 mmol). The reaction mixture was stirred overnight at room temperature. The product mixture was concentrated under vacuum. The residue was purified using column chromatography on silica gel eluting with 50 % -

100 % ethyl acetate in hexanes. Collection and concentration of appropriate fractions provided the title compound as pale yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 7.25 (m, 2H), 7.01 (t, *J* = 8.7 Hz, 2H), 4.64 (d, *J* = 14.6 Hz, 1H), 4.48 (d, *J* = 14.5 Hz, 1H), 4.34 (m, 2H), 3.27 (m, 2H), 2.23 (dd, *J* = 17, 10.6 Hz, 1H), 2.0 (m, 1H), 1.61 (m, 1H).

5

Step 5: 4-(Aminomethyl)-1-(4-fluorobenzyl)piperidin-2-one.

To a degassed solution of 1-(4-fluorobenzyl)-4-(nitromethyl)piperidin-2-one (1.4 g, 5.3 mmol) in absolute ethanol (80 mL) was added wet Raney nickel catalyst (1.4 g of a 50 % weight slurry in water) which was washed to neutral pH with deionized water and absolute ethanol. The reaction mixture was shaken under an atmosphere of 55 psi of hydrogen for 16 hours. The product mixture was filtered through celite and the filtrate was concentrated under vacuum to afford the title compound as colorless viscous oil. ¹H NMR (400 MHz, CDCl₃) δ 7.24 (m, 2H), 7.01 (t, *J* = 8.7 Hz, 2H), 4.64 (d, *J* = 14.6 Hz, 1H), 4.44 (d, 1H), 3.23 (m, 2H), 2.65 (m, 2H), 2.16- 1.28 (m).

15 Step 6: Ethyl ([1-(4-fluorobenzyl)-2-oxopiperidin-4-yl]methyl)amino)-(oxo)acetate.

To a cooled (0 °C) solution of 4-(aminomethyl)-1-(4-fluorobenzyl)-piperidin-2-one (1.20 g, 5.07 mmol) in anhydrous methylene chloride (20 mL) under an atmosphere of nitrogen was added diisopropylethylamine (0.72 g, 5.58 mmol). Ethyl oxalyl chloride (0.69 g, 5.07 mmol) was added to the reaction mixture over seven minutes. The reaction stirred at 0 °C for 1 hour, then warmed to room temperature for 2 hours. The resultant solution was diluted with ice water and methylene chloride (100 mL). The organic extract was dried over Na₂SO₄, filtered, and concentrated under vacuum. The residual oil was purified using column chromatography on silica gel eluting with 20 % - 80 % ethyl acetate in hexanes. Collection and concentration of appropriate fractions provided the title compound as a colorless viscous oil. ¹H NMR (400 MHz, CDCl₃) δ 7.22 (m, 3H), 7.00 (t, *J* = 8.6 Hz, 2H), 4.60 (d, *J* = 14.5 Hz, 1H), 4.49 (d, *J* = 14.5 Hz, 1H), 4.35 (q, *J* = 7.1 Hz, 2H), 3.34-3.15 (m, 4H), 2.61 (m, 1H), 2.15 (m, 2H), 1.92 (m, 1H), 1.51 (m, 1H), 1.39 (t, *J* = 7.1 Hz, 1H).


Step 7: 2-(4-Fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione.

30 To a cooled (0 °C) solution of rigorously dried ethyl ([1-(4-fluorobenzyl)-2-oxopiperidin-4-yl]methyl)amino)-(oxo)acetate (0.273 g, 0.812 mmol) in anhydrous DMF (1.5 mL) under an atmosphere of nitrogen, was added a solution of LiHMDS in THF (0.85 M, 1 M; 0.85 mmol). The reaction mixture stirred for 30 min at 0 °C. Methyl iodide (0.115 g, 0.812 mmol) was added and the mixture was stirred for 2 hours at 0 °C, then at room temperature for 16 hours. The reaction was then

cooled to 0 °C and treated with a solution of LiHMDS in THF (0.85 mL, 1 M; 0.85 mmol) and stirred for 1 hour at 0 °C, then for 20 hours at room temperature. The product mixture was concentrated under vacuum and purified using HPLC on C18 stationary phase eluting with water/acetonitrile/TFA mobile phase. Collection and lyophilization of the appropriate fractions provided the title compound as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 13.66 (br s, 1H), 7.23 (m, 2H), 7.09 (t, *J* = 8.7 Hz, 2H), 4.74 (d, *J* = 14.7 Hz, 1H), 4.46 (d, *J* = 14.7 Hz, 1H), 3.31 (m, 4H), 3.06 (s, 3H), 3.03 (m, 1H), 1.89 (dq, *J* = 13, 3.3 Hz, 1H), 1.55 (qd, *J* = 13, 5.2 Hz, 1H).

EXAMPLE 2

2-(4-Fluorobenzyl)-8-hydroxy-5,5,6-trimethyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione



Step 1: 1-(4-Fluorobenzyl)-4-(1-methyl-1-nitroethyl)piperidin-2-one.

To a solution 1-(4-fluorobenzyl)-5,6-dihydropyridin-2(1H)-one (see Step 3 of Example 1) (0.75 g, 3.65 mmol) in 2-nitropropane (9.77 g, 109 mmol) was added DBU (0.56 g, 3.65 mmol). After stirring for 48 hours, the reaction mixture was concentrated under vacuum. The residual material was purified using silica gel column chromatography eluting with CH₂Cl₂. The appropriate fractions were combined and concentrated to afford the title compound. ¹H NMR (400 MHz, CDCl₃) δ 7.44 (m, 2H), 7.04 (t, *J*=9 Hz, 2H), 4.64 (d, *J*=15 Hz, 1H), 4.43 (d, *J*=15 Hz, 1H), 3.23 (m, 2H), 2.56 (m, 2H), 2.22 (m, 1H), 1.73 (m, 1H), 1.56 (s, 6H), 1.51 (m, 1H) ppm. ES MS *M*+1 = 295.

Step 2: 4-(1-Amino-1-methylethyl)-1-(4-fluorobenzyl)piperidine-2-one.

To a solution of 1-(4-fluorobenzyl)-4-(1-methyl-1-nitroethyl)piperidin-2-one (1.05 g, 3.57 mmol) in degassed ethanol (75 mL) was added Raney Nickel (2 g of a 50% by weight slurry in water). The reaction mixture was shaken under 55 psi of hydrogen for 4 hours. The reaction mixture was filtered through a bed of celite, then concentrated under vacuum to give an oil that solidified on standing. The solid was triturated in ether and collected by filtration to give the title compound. ¹H NMR (400 MHz, CDCl₃) δ 7.22 (m, 2H), 7.02 (t, *J*=8.6 Hz, 2H), 4.65 (d, *J*=14.5 Hz, 1H), 4.44 (d, *J*=14.6 Hz, 1H), 3.26 (m, 2H), 2.60 (d, *J*=5 Hz, 1H), 2.25 (m, 1H), 1.97 (d, *J*=11.5 Hz, 1H), 1.75 (m, 1H), 1.44 (m, 1H), 1.11 (d, *J*=9 Hz, 6H) ppm. ES MS *M*+1 = 265.

Step 3: Ethyl ({1-[1-(4-fluorobenzyl)-2-oxopiperidin-4-yl]-1-methylethyl}amino)(oxo)acetate.

To a cooled (0 °C) solution of 4-(1-amino-1-methylethyl)-1-(4-fluorobenzyl)piperidine-2-one (0.92 g, 3.5 mmol) and diisopropylethylamine (0.49 g, 3.8 mmol) in CH₂Cl₂ (10 mL) was added ethyl oxalyl chloride (0.48 g, 3.5 mmol). After stirring at 0 °C for 1 hour, the reaction mixture was partitioned between CH₂Cl₂ and water. The organic extract was dried with Na₂SO₄, filtered, and concentrated under vacuum to provide the title compound as an oil. ¹H NMR (400 MHz, CDCl₃) δ 7.22 (m, 2H), 7.01 (m, 2H), 6.93 (br, 1H), 4.59 (d, *J*=12 Hz, 1H), 4.49 (d, *J*=12 Hz, 1H), 3.34 (m, 2H), 3.22 (m, 2H), 2.70 (m, 1H), 2.54 (m, 1H), 2.19 (m, 1H), 1.87 (m, 1H), 1.47 (m, 1H), 1.18 (m, 7H), 1.14 (m, 1H) ppm. ES MS *M*+1 = 365.


Step 4: 2-(4-Fluorobenzyl)-8-hydroxy-5,5,6-trimethyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione.

To a cooled (0 °C) solution of ethyl ({1-[1-(4-fluorobenzyl)-2-oxopiperidin-4-yl]-1-methylethyl}amino)(oxo)acetate (1.2 g, 3.3 mmol) in anhydrous DMF (10 mL) under an atmosphere of nitrogen was added LiHMDS (1M in THF, 3.5 mL, 3.5 mmol). After stirring at 0 °C for 15 min, methyl iodide (0.47 g, 3.3 mmol) was added to the reaction mixture. After stirring for an additional 15 min, more LiHMDS (1M in THF, 3.5 mmol, 3.5 mmol) was added. After another 2h, an additional portion of LiHMDS (1M in THF, 3.5 mL, 3.5 mmol) was added. After stirring at room temperature for 24 hours, the reaction mixture was concentrated under vacuum. The residual material was purified using reverse phase HPLC on a C18 stationary phase eluting with a gradient of 5 % - 95 % acetonitrile (0.1 % TFA) in H₂O (0.1 % TFA) to afford the title compound as an oil. ¹H NMR (400 MHz, CDCl₃) δ 7.27 (m, 2H), 7.04 (t, *J*=8.6 Hz, 2H), 4.67 (d, *J*=14.7 Hz, 1H), 4.52 (d, *J*=14.7 Hz, 1H), 3.33 (d, *J*=6 Hz, 2H), 3.31 (s, 3H), 2.84 (m, 1H), 1.93 (m, 1H), 1.64 (m, 1H), 1.37 (s, 3H), 1.13 (s, 3H) ppm. ES MS *M*+1 = 333.

The enantiomers of the title compound were obtained by separation using a ChiralPak AD column with 100% methanol as the mobile phase. On an analytical ChiralPak AD column eluting with 100% methanol, the first enantiomer had a retention time of 5.05 min and a negative sign of rotation, the second enantiomer had a retention time of 7.07 min and a positive sign of rotation.

EXAMPLE 3

2-(4-Fluorobenzyl)-8-hydroxy-6-methyl-4a-phenyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione



Step 1: 4-Cyano-4-phenyl-1-butene.

5 To a cooled solution (-78 °C) of benzyl cyanide (5.00 g, 42.7 mmol) in anhydrous THF was added LiHMDS (1M in THF, 46.9 mL, 46.9 mmol) dropwise. After 0.5 hour, allyl bromide (5.16 g, 42.6 mmol) was added dropwise to the reaction mixture. The reaction mixture was warmed to room temperature, and the solvent was removed under vacuum. The residual material was purified using silica gel column chromatography eluting with 100% hexanes. The appropriate fractions were combined and concentrated to afford the title compound. ¹H NMR (400 MHz, DMSO-d₆) δ 7.44 (m, 5H), 5.81 (m, 1H), 5.12 (m, 2H), 4.35 (t, J=6.9 Hz, 1H), 2.61 (m, 2H) ppm. ES MS M+1 = 158.

Step 2: Methyl 3-cyano-3-phenylhex-5-enoate.

15 To a cooled solution (-78 °C) of 4-cyano-4-phenyl-1-butene (6.23 g, 39.6 mmol) in anhydrous THF was added LiHMDS (1M in THF, 43.5 mL, 43.5 mmol) dropwise. The solution was stirred at -78 °C for 0.5 hour, then methyl bromoacetate (6.67 g, 43.6 mmol) was added dropwise. After stirring at room temperature for 24 hours, the reaction mixture was concentrated under vacuum. The residual material was purified using silica gel column chromatography eluting with 0 - 20% EtOAc in hexanes. The appropriate fractions were combined and concentrated to afford the title compound. ¹H NMR (400 MHz, DMSO-d₆) δ 7.38 (m, 5H), 5.57 (m, 1H), 5.14 (m, 2H), 3.49 (s, 3H), 3.49 (m, 2H), 2.78 (m, 2H) ppm. ES MS M+1 = 230.

Step 3: 3-Cyano-3-phenylhex-5-enoic acid.

25 To a solution of methyl 3-cyano-3-phenylhex-5-enoate (6.03 g, 26.3 mmol) in methanol (90 mL) was added 1M aqueous NaOH (29 mL, 29 mmol). After stirring overnight at room temperature, the reaction mixture was concentrated under vacuum. The residue was partitioned between EtOAc and water and the pH of the aqueous phase was adjust to pH 4 using 3N aqueous HCl. The organic extract was dried with Na₂SO₄, filtered, and concentrated under vacuum to provide the title compound as a

yellow oil. ^1H NMR (400 MHz, DMSO- d_6) δ 12.45 (br, 1H), 7.50 (d, $J=1.6$ Hz, 2H), 7.45 (t, $J=1.9$ Hz, 2H), 7.42 (t, $J=1.7$ Hz, 1H), 5.51 (m, 1H), 5.11 (m, 2H), 3.17 (s, 2H), 2.77 (m, 2H), 2.51 (s, 2H) ppm. ES MS $M+1 = 216$.

5 Step 4: 3-Cyano-*N*-(4-fluorobenzyl)-3-phenylhex-5-enamide.

To a solution of 3-cyano-3-phenylhex-5-enoic acid (5.66 g, 26.3 mmol) and 4-fluorobenzylamine (3.95 g, 31.6 mmol) in DMF was added 1-hydroxybenzotriazole hydrate (6.04 g, 39.4 mmol), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (7.56 g, 39.4 mmol), and triethylamine (3.19 g, 31.5 mmol). After stirring overnight at room temperature, the reaction mixture was concentrated under vacuum. The residue was partitioned between EtOAc and water. The organic extract was dried with Na_2SO_4 , filtered, and concentrated under vacuum to provide the title compound as a light yellow solid. ^1H NMR (400 MHz, DMSO- d_6) δ 8.41 (s, 1H), 7.47 (d, $J=7.9$ Hz, 2H), 7.36 (t, $J=7$ Hz, 2H), 7.35 (t, $J=7$ Hz, 1H), 7.04 (t, $J=8.9$ Hz, 2H), 6.97 (t, $J=8.6$ Hz, 2H), 5.53 (m, 1H), 5.12 (m, 2H), 4.21 (m, 1H), 4.08 (m, 1H), 2.88 (m, 3H), 2.77 (m, 1H) ppm. ES MS $M+1 = 323$.

15

Step 5: 4-Cyano-1-(4-fluorobenzyl)-4-phenyl-1,2,3,4-tetrahydropyridin-2-one.

Into a cooled (-78°C) solution of 3-cyano-*N*-(4-fluorobenzyl)-3-phenylhex-5-enamide (8.00 g, 24.8 mmol) in CH_2Cl_2 (500 mL) was bubbled ozone until a blue color persisted. The mixture was stirred for 10 min, then dimethyl sulfide (84.6 g, 1.36 mol) was added. After stirring for 48 hours at room temperature, the mixture was concentrated under vacuum. The residue was partitioned between EtOAc and water. The organic extract was dried with Na_2SO_4 , filtered, and concentrated under vacuum. The residual material was purified using silica gel column chromatography eluting with 0 - 40% EtOAc in hexanes. The appropriate fractions were combined and concentrated to afford the title compound. ^1H NMR (400 MHz, DMSO- d_6) δ 7.45 (m, 5H), 7.29 (m, 2H), 7.15 (t, $J=8.9$ Hz, 2H), 6.78 (d, $J=7.7$ Hz, 1H), 5.50 (d, $J=7.7$ Hz, 1H), 4.72 (s, 2H), 3.17 (q, $J=16$ Hz, 2H) ppm. ES MS $M+1 = 307$.

25

Step 6: 4-Cyano-1-(4-fluorobenzyl)-4-phenylpiperidin-2-one.

To a solution of 4-cyano-1-(4-fluorobenzyl)-4-phenyl-1,2,3,4-tetrahydropyridin-2-one (3.98 g, 13.0 mmol) in degassed methanol was added 10% Pd/C (0.8 g). The reaction shaken under an atmosphere of 33 psi of hydrogen for 18 hours. More 10% Pd/C (0.4 g) was added and the mixture was shaken under 45 psi of hydrogen for 48 hours. The mixture was filtered through a bed of celite and the filtrate was concentrated to give the title compound as a white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 7.55 (d, $J=7.7$ Hz, 2H), 7.46 (t, $J=7.9$ Hz, 2H), 7.38 (t, $J=7.3$ Hz, 1H), 7.28 (t, $J=3$ Hz, 2H), 7.16 (t, $J=2.4$

30

Hz, 2H), 4.58 (s, 2H), 3.44 (m, 1H), 3.25 (m, 1H), 3.07 (q, $J=16$ Hz, 2H), 2.41 (s, 2H) ppm. ES MS $M+1 = 309$.

Step 7: 1-(4-Fluorobenzyl)-4-phenylpiperidin-2-one-4-carbothioamide.

5 To a cooled (0 °C) solution of 4-cyano-1-(4-fluorobenzyl)-4-phenylpiperidin-2-one (3.39 g, 10.9 mmol) in pyridine (77 mL) was added triethylamine (45.0 g, 0.77 mol). The reaction mixture was saturated with hydrogen sulfide gas, the reaction vessel was sealed and then brought to ambient temperature. After stirring for 24 hours, the reaction mixture was cooled (0 °C), the reaction vessel was opened, then the solvents were concentrated under vacuum. The residual material was purified using
10 silica gel column chromatography eluting with 0 - 70% EtOAc in hexanes. The appropriate fractions were combined and concentrated to afford the title compound. ^1H NMR (400 MHz, DMSO- d_6) δ 9.65 (s, 1H), 8.68 (s, 1H), 7.35 (m, 4H), 7.26 (t, $J=7$ Hz, 1H), 7.13 (t, $J=8.3$ Hz, 2H), 7.04 (t, $J=8.9$ Hz, 2H), 4.48 (d, $J=15$ Hz, 1H), 4.35 (d, $J=15$ Hz, 1H), 3.16 (m, 2H), 3.93 (d, $J=17$ Hz, 1H), 2.82 (m, 1H), 2.51 (m, 2H) ppm. ES MS $M+1 = 343$.

15 Step 8: 4-Aminomethyl-1-(4-fluorobenzyl)-4-phenylpiperidin-2-one.
To a solution of Raney nickel (50% by weight slurry in water, 0.66 g, 11.3 mmol) in ethanol was added sodium hydroxide (0.45 g, 11.3 mmol) and the mixture was heated to 50 °C under an atmosphere of nitrogen. After 0.5 hour, 1-(4-fluorobenzyl)-4-phenylpiperidin-2-one-4-carbothioamide
20 (1.94 g, 5.66 mmol) in ethanol (5 mL) was added and the reaction mixture was stirred at 50 °C for 1.5 hours. The solids were then removed by filtration through celite and the filtrate was concentrated under vacuum. The residual material was purified using reverse phase HPLC on a C18 stationary phase eluting with a gradient of 5 % - 95 % acetonitrile (0.1 % TFA) in H_2O (0.1 % TFA) to afford the TFA salt of the title compound as a white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 7.61 (br, 2H), 7.39 (m, 5H), 6.97 (m,
25 4H), 4.61 (d, $J=15$ Hz, 1H), 4.13 (d, $J=15$ Hz, 1H), 3.12 (m, 4H), 2.67 (d, $J=7$ Hz, 1H), 2.51 (m, 1H), 2.23 (m, 1H), 2.18 (m, 1H) ppm. ES MS $M+1 = 313$.

Step 9: Ethyl ({[1-(4-fluorobenzyl)-2-oxo-4-phenylpiperidin-4-yl]methyl}amino)(oxo)acetate.

30 To a cooled (0 °C) solution of 4-(aminomethyl)-1-(4-fluorobenzyl)-4-phenylpiperidin-2-one (0.40 g, 1.28 mmol) in CH_2Cl_2 was added diisopropylethylamine (0.19 g, 1.53 mmol) and ethyl oxalyl chloride (0.19 g, 1.41 mmol). After 1 hour, the reaction mixture was concentrated. The residual material was purified using reverse phase HPLC on a C18 stationary phase eluting with a gradient of 5 % - 95 % acetonitrile (0.1 % TFA) in H_2O (0.1 % TFA) to afford the title compound as a white solid. ^1H NMR (400 MHz, DMSO- d_6) δ 8.67 (m, 1H), 7.34 (m, 5H), 6.95 (m, 4H), 4.58 (d, $J=15$ Hz, 1H), 4.21 (q,


$J=7$ Hz, 2H), 4.11 (d, $J=15$ Hz, 1H), 3.38 (m, 1H), 3.28 (m, 1H), 3.09 (m, 1H), 2.83 (d, $J=17$ Hz, 1H), 2.56 (d, $J=13$ Hz, 2H), 2.21 (m, 1H), 1.98 (m, 1H), 1.25 (t, $J=7$ Hz, 3H) ppm. ES MS $M+1 = 413$.

Step 10: 2-(4-Fluorobenzyl)-8-hydroxy-6-methyl-4a-phenyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione.

To a cooled (0 °C) solution of ethyl ({[1-fluorobenzyl]-2-oxo-4-phenylpiperidin-4-yl}methyl)amino(oxo)acetate (0.27 g, 0.65 mmol) in DMF was added LiHMDS (0.71 mL of a 1.0 M solution in THF, 0.71 mmol) and iodomethane (0.34 g, 2.01 mmol). After stirring at room temperature under anhydrous conditions for 6 hours, the reaction was cooled to 0 °C and more LiHMDS (1.94 mmol) was added. After stirring overnight at room temperature, the reaction mixture was filtered and concentrated. The residual material was purified using reverse phase HPLC on a C18 stationary phase eluting with a gradient of 5 % - 95 % acetonitrile (0.1 % TFA) in H₂O (0.1 % TFA) to give a solid which was crystallized from methanol to afford the title compound. ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.31 (m, 7H), 7.15 (m, 2H), 4.66 (d, $J=3.3$ Hz, 1H), 4.63 (d, $J=3.2$ Hz, 1H), 3.85 (d, $J=3.7$ Hz, 1H), 3.82 (d, $J=3.5$ Hz, 1H), 3.17 (d, $J=12$ Hz, 1H), 2.68 (s, 3H), 2.55 (m, 1H), 2.16 (d, $J=11$ Hz, 1H), 1.98 (m, 1H) ES MS $M+1 = 381$. The enantiomers of the title compound were separated using a ChiralPak AD column with 50% methanol in ethanol as the mobile phase. On an analytical ChiralPak AD column eluting with 50% methanol in ethanol, the first enantiomer had a retention time of 5.92 min and a negative sign of rotation, the second enantiomer had a retention time of 8.98 min and a positive sign of rotation.

EXAMPLE 4

5-(*tert*-Butyloxycarbonyl)-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione



Step 1: *tert*-Butyl *N*-[ethoxy(oxo)acetyl]-*N*-methylglycinate.

To a solution *tert*-butyl *N*-methylglycinate hydrochloride (1.82 g, 10.0 mmol) in 1,2-dichloroethane (15 mL) was added triethylamine (2.13 g, 21.0 mmol). After stirring for 0.5 hour at room temperature, the reaction mixture was cooled to 0 °C and ethyl oxalyl chloride (1.44 g, 10.5 mmol) was


added dropwise over 10 min. After warming to room temperature and stirring for 24 hours, the reaction mixture was partitioned between 1, 2-dichloroethane and water. The organic extract was dried with Na_2SO_4 , filtered, and concentrated under vacuum to provide the title compound as a yellow oil. ^1H NMR (300 MHz, CDCl_3) δ 4.32 (m, 2H), 4.03 (d, $J=3.9$ Hz, 2H), 3.06 (d, $J=10.3$ Hz, 3H), 1.48 (s, 9H), 1.38 (m, 3H) ppm. ES MS $M+1 = 246$.

Step 2: 5-(tert-Butyloxycarbonyl)-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione.

To a cooled (-78°C) solution of *tert*-butyl *N*-[ethoxy(oxo)acetyl]-*N*-methylglycinate (1.43 g, 5.85 mmol) and 1-(4-fluorobenzyl)-5,6-dihydropyridin-2(1H)-one from Step 3 of Example 1 (1 g, 5 mmol) in anhydrous THF (10 mL) was added LiHMDS (1 M in THF) (1.02 g, 6.09 mmol) dropwise. The reaction mixture was stirred for 10 min. at -78°C , warmed to room temperature for 2.5 hours, then heated to 40°C . After heating for 24 hours, the reaction mixture was quenched with cold diluted HCl, then diluted with methanol, and concentrated under vacuum. The residual material was purified using reverse phase HPLC on a C18 stationary phase eluting with 5 % - 95 % acetonitrile (0.1 % TFA) in H_2O (0.1 % TFA) to give the title compound as a mixture of diastereomers. The appropriate fractions were collected for the first-eluting diastereomer, diastereomer 1, and this material was purified further using reverse phase HPLC on a C18 stationary phase eluting with 15 % -85 % acetonitrile (0.1 % TFA) in H_2O (0.1 % TFA). For the second eluting diastereomer, diastereomer 2, the appropriate fractions were collected and the material was purified further using reverse phase HPLC on a C18 stationary phase eluting with 10 %-60% methanol (0.1 % TFA) in H_2O (0.1 % TFA). Diastereomer 1 : (4a*R*, 5*R*)-5-(tert-Butyloxycarbonyl)-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione. ^1H NMR (400 MHz, CDCl_3) δ 6.84 (m, 2H), 6.62 (t, $J=8.7$ Hz, 2H), 4.31 (d, $J=14.7$ Hz, 1H), 4.05 (d, $J=14.7$ Hz, 1H), 3.43 (d, $J=6.5$ Hz, 1H), 2.91 (m, 2H), 2.69 (m, 1H), 2.57 (s, 3H), 1.42 (m, 1H), 1.21 (m, 1H), 1.15 (s, 9H) ppm. ES MS $M+1 = 405$. Diastereomer 2 : (4a*R*, 5*S*)-5-(tert-Butyloxycarbonyl)-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione. ^1H NMR (400 MHz, CDCl_3) δ 13.07 (s, 1H), 7.22 (m, 2H), 7.02 (t, $J=8.7$ Hz, 2H), 4.67 (d, $J=14.7$ Hz, 1H), 4.51 (d, $J=14.7$ Hz, 1H), 3.82 (d, $J=6.5$ Hz, 1H), 3.33 (m, 1H), 2.99 (s, 3H), 2.02 (m, 3H), 1.68 (m, 1H), 1.44 (s, 9H) ppm. ES MS $M+1 = 405$.

EXAMPLE 5

5-Ethyl-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione



Step 1: 1-(4-Fluorobenzyl)-4-(1-nitropropyl)piperidin-2-one.

To a solution 1-(4-fluorobenzyl)-5,6-dihydropyridin-2(1H)-one (2.0 g, 10 mmol) in 1-nitropropane (8.68 g, 97.5 mmol) was added DBU (1.48 g, 9.75 mmol). After stirring for 24 hours, the reaction mixture was concentrated under vacuum. The residual material was purified by silica gel column chromatography eluting with 1% methanol in CH_2Cl_2 . The appropriate fractions were combined and concentrated to afford the title compound as mixture of diastereomers. The diastereomers co-eluted with a retention time of 3.01 min. The HPLC method was performed on a Hewlett-Packard Zorbax SB-C8 column (75 x 4.6 mm, 3.5 micron) eluting with 5 %– 100 % CH_3CN (0.1% TFA) in H_2O (0.1% TFA). ES MS $M+1 = 295$.

Step 2: 4-(1-Aminopropyl)-1-(4-fluorobenzyl)piperidin-2-one.

To a solution of 1-(4-fluorobenzyl)-4-(1-nitropropyl)piperidin-2-one (2.8 g, 9.51 mmol) in ethanol (50 mL) was added Raney Nickel (2 grams of a 50% by weight slurry in water). The reaction mixture was shaken under 55 psi of hydrogen. After 24 hours, the reaction mixture was filtered through a bed a celite, then concentrated under vacuum to afford the title compound as a mixture of diastereomers. The first diastereomer has a retention time of 2.05 min and the second diastereomer has a retention time of 2.10 min. The HPLC method was performed on a Hewlett-Packard Zorbax SB-C8 column (75 x 4.6 mm, 3.5 micron) eluting with 5 %– 100 % CH_3CN (0.1% TFA) in H_2O (0.1% TFA). ES MS $M+1 = 265$.


Step 3: 5-Ethyl-2-(4-fluorobenzyl)-8-hydroxy-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione.

To a cooled (-78°C) solution of 4-(1-aminopropyl)-1-(4-fluorobenzyl)piperidin-2-one (2.0 g, 8 mmol) in anhydrous THF (20 mL) was added LiHMDS (1M in THF, 18.9 mL, 18.9 mmol) under an atmosphere of nitrogen. After stirring at -78°C for 5 min, diethyl oxalate (3.32 g, 22.7 mmol) was added to the reaction mixture. After stirring for 2 hours at -78°C , then for 0.5 hour at -20°C ,

LiHMDS (1M in THF, 22 mL, 22 mmol) was added. After stirring at room temperature for 24 hours, the reaction mixture was quenched with 1N HCl. The mixture was partitioned between EtOAc and water. The organic extract was dried with Na₂SO₄, filtered, concentrated under vacuum, then filtered through a plug of silica gel and eluting with 1% methanol in CH₂Cl₂. The diastereomers were separated using
 5 reverse phase HPLC on a C18 stationary phase eluting with 5 % - 95 % acetonitrile (0.1 % TFA) in H₂O (0.1 % TFA) to afford the title compounds. Diastereomer 1: ¹H NMR (400 MHz, CDCl₃) δ 13.67 (br, 1H), 7.26 (m, 2H), 7.04 (t, *J* = 8.6 Hz, 2H), 4.68 (d, *J* = 14.7 Hz, 1H), 4.52 (d, *J* = 14.7 Hz, 1H), 3.35 (m, 4H), 1.79 (m, 3H), 1.48 (m, 1H), 0.97 (t, *J* = 7.3 Hz, 3H) ppm. ES MS *M*+1 = 319. Diastereomer 2 : ¹H NMR (400 MHz, CDCl₃) δ 7.26 (m, 2H), 7.04 (t, *J* = 8.6 Hz, 2H), 6.54 (br, 1H), 4.72 (d, *J* = 14.7 Hz, 1H), 4.49 (d, *J* = 14.7 Hz, 1H), 3.34 (m, 3H), 2.73 (dt, *J* = 4, 13 Hz, 1H), 1.99 (m, 1H), 1.78 (m, 1H), 1.57 (m, 2H), 1.01 (t, *J* = 7.5 Hz, 3H) ppm. ES MS *M*+1 = 319.

EXAMPLE 6

(4a*R*, 5*R*)-5-Ethyl-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione
 15 dione



Step 1: 5-Ethyl-2-(4-fluorobenzyl)-8-methoxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione.


To a solution of 5-ethyl-2-(4-fluorobenzyl)-8-hydroxy-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione, Diastereomer 1 from Example 5 (0.46 g, 1.5 mmol) in THF (30 mL) was added cesium carbonate (1.4 g, 4.3 mmol) and methyl iodide (1.02 g, 7.19 mmol). After stirring for 24 hours at room temperature, the reaction mixture was partitioned between EtOAc and 1N HCl. The organic extract was dried with Na₂SO₄, filtered, and concentrated under vacuum. The residual material was purified using reverse phase HPLC on a C18 stationary phase eluting with 5 % - 95 % acetonitrile (0.1 % TFA) in H₂O (0.1 % TFA) to afford the title compound as a yellow oil. ¹H NMR (400 MHz, CDCl₃) δ 7.51 (m, 2H), 7.05 (t, *J* = 9 Hz, 2H), 5.12 (d, *J* = 15 Hz, 1H), 4.73 (d, *J* = 15 Hz, 1H), 4.56 (d, *J* = 15 Hz, 1H), 4.17 (d, *J* = 15 Hz, 1H), 3.28 (m, 3H), 3.13 (d, *J* = 15 Hz, 3H), 2.39 (m, 1H), 2.18 (m, 1H), 1.67 (m, 4H), 0.98 (s, 3H) ppm. ES MS *M*+1 = 347.

Step 2: 5-Ethyl-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione.

To a cooled (0 °C) solution of 5-ethyl-2-(4-fluorobenzyl)-8-methoxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione (0.26 g, 0.75 mmol) in CH₂Cl₂ (20 mL) was added boron tribromide (0.94 g, 3.8 mmol). After stirring for 0.5 hour at 0 °C, the reaction mixture was quenched with methanol, then concentrated under vacuum. The residual material was purified using reverse phase HPLC on a C18 stationary phase eluting with 5 % - 95 % acetonitrile (0.1 % TFA) in H₂O (0.1 % TFA) to afford the title compound. ¹H NMR (400 MHz, CDCl₃) δ 7.24 (m, 2H), 7.05 (t, *J*=9 Hz, 2H), 5.12 (d, *J*=14.6 Hz, 1H), 4.73 (d, *J*=14.7 Hz, 1H), 3.14 (s, 3H), 1.76 (m, 4H), 0.98 (t, *J*=7.5 Hz, 3H) ppm. ES MS *M*+1 =333.

EXAMPLE 7

(4a*R*, 5*S*)-5-Ethyl-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione



Step 1: 5-Ethyl-2-(4-fluorobenzyl)-8-methoxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione.

The title compound was prepared from 5-ethyl-2-(4-fluorobenzyl)-8-hydroxy-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione, Diastereomer 2 from Example 5, using the procedure given in Step 1 of Example 6. ES MS *M*+1 = 347.


Step 2: 5-Ethyl-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione.

To a cooled (0 °C) solution of 5-ethyl-2-(4-fluorobenzyl)-8-methoxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione, (0.39 g, 1.1 mmol) in CH₂Cl₂ (20 mL) was added hydrogen bromide (30 % in HOAc, 5 mL). After stirring for 3 hours at 0 °C, the reaction mixture was concentrated under vacuum. The residual material was purified using reverse phase HPLC on a C18 stationary phase eluting with 5 % - 95 % acetonitrile (0.1 % TFA) in H₂O (0.1 % TFA) to afford the title compound. ¹H NMR (400 MHz, CDCl₃) δ 13.73 (s, 1H), 7.26 (m, 2H), 7.04 (t, *J*=9 Hz, 2H), 4.69 (d,

$J=14.6$ Hz, 1H), 4.49 (d, $J=14.7$ Hz, 1H), 3.36 (m, 3H), 3.04 (s, 3H), 2.83 (dt, $J=12.3$; 4.2 Hz, 1H), 1.97 (m, 2H), 1.68 (m, 2H), 0.92 (t, $J=7.3$ Hz, 3H) ppm. ES MS $M+1=333$.

EXAMPLE 8

- 5 6-(Cyclopropylmethyl)-2-(4-fluorobenzyl)-8-hydroxy-5,5-dimethyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione



Step 1: 4-{1-[(Cyclopropylmethyl)amino]-1-methylethyl}-1-(4-fluorobenzyl)piperidin-2-one.

To a suspension of 4-(1-amino-1-methylethyl)-1-(4-fluorobenzyl)piperidin-2-one from

- 10 Example 2, Step 2, (1.00 g, 3.78 mmol) in acetonitrile (20 mL) was added cesium carbonate (2.46 g, 7.56 mmol) and bromomethylcyclopropane (1.02 g, 7.56 mmol). After stirring at room temperature for 48 hours, more bromomethylcyclopropane (0.36 g, 2.67 mmol) was added. After 24 hours, more bromomethylcyclopropane (0.36 g, 2.67 mol) was added. After another 24 hours, the product mixture was concentrated. The residue was partitioned between ethyl acetate and H_2O . The organic extract was
- 15 washed with brine, dried with Na_2SO_4 , filtered, and concentrated under vacuum. The residual material was purified using reverse phase HPLC on a C18 stationary phase eluting with a gradient of 5 % - 95 % acetonitrile (0.1 % TFA) in H_2O (0.1 % TFA) to afford the title compound as the TFA salt. The TFA salt was dissolved in saturated $NaHCO_3$ and extracted with EtOAc. The combined organic extract was dried over Na_2SO_4 , filtered, and concentrated under vacuum to give the free base of the title compound. 1H
- 20 NMR (400 MHz, $CDCl_3$) δ 7.21 (m, 2H), 7.01 (t, $J=10$ Hz, 2H), 4.64 (d, $J=16$ Hz, 1H), 4.46 (d, $J=16$ Hz, 1H), 3.24 (m, 1H), 3.15 (m, 1H), 2.55 (m, 1H), 2.34 (d, $J=6.8$ Hz, 2H), 2.24 (m, 1H), 1.94 (m, 1H), 1.81 (m, 1H), 1.46 (m, 1H), 1.01 (s, 6H), 0.86 (m, 1H), 0.45 (d, $J=1.1$ Hz, 2H), 0.082 (t, $J=5.9$ Hz, 2H) ppm. ES MS $M+1=319$.

- 25 Step 2: Ethyl ((cyclopropylmethyl){1-[1-(4-fluorobenzyl)-2-oxopiperidin-4-yl]-1-methylethyl}amino)(oxo)acetate.

- To a solution of 4-{1-[(cyclopropylmethyl)amino]-1-methylethyl}-1-(4-fluorobenzyl)piperidin-2-one (1.4 g, 4.4 mmol) in CH_2Cl_2 (15 mL) was added DIEA (0.625 g, 4.83 mmol) and ethyl oxalyl chloride (0.6 g, 4.4 mmol). After 1.5 hours, the reaction mixture was partitioned
- 30 between CH_2Cl_2 and water. The organic extract was washed with brine, dried with Na_2SO_4 , filtered,


and concentrated under vacuum. The residue was purified by silica gel column chromatography eluting with 1 - 3 % methanol in CH₂Cl₂. The appropriate fractions were combined and concentrated to afford the title compound. ¹H NMR (400 MHz, CDCl₃) δ 7.22 (m, 2H), 7.01 (t, *J*=8 Hz, 2H), 4.68 (d, *J*=16 Hz, 1H), 4.38 (d, *J*=16 Hz, 1H), 4.30 (q, *J*=8 Hz, 2H), 3.34 (m, 1H), 3.19 (m, 4H), 2.56 (m, 1H), 2.21 (m, 1H), 1.82 (m, 1H), 1.49 (s, 3H), 1.44 (m, 1H), 1.41 (s, 3H), 1.35 (q, *J*=7 Hz, 3H), 1.01 (m, 1H), 0.61 (m, 2H), 0.25 (m, 2H) ppm. ES MS *M*+1 = 419.

Step 3: 6-(Cyclopropylmethyl)-2-(4-fluorobenzyl)-8-hydroxy-5,5-dimethyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione.

To a cooled solution (0 °C) of ethyl ((cyclopropylmethyl){1-[1-(4-fluorobenzyl)-2-oxopiperidin-4-yl]-1-methylethyl}amino)(oxo)acetate (1.56 g, 3.73 mmol) in DMF (15 mL) was added LiHMDS (1M in THF, 5.59 mL, 5.59 mmol) dropwise over 1 min. After stirring for 3 hours, the product mixture was concentrated under vacuum and purified using reverse phase HPLC on C18 stationary phase eluting with a gradient of 5 % - 95 % acetonitrile (0.1 % TFA) in H₂O (0.1 % TFA) to afford the title compound as a pale yellow solid. ¹H NMR (400 MHz, CDCl₃) δ 7.26 (m, 2H), 7.04 (t, *J*=8.6 Hz, 2H), 4.61 (s, 2H), 3.41 (d, *J*=6.8 Hz, 2H), 3.32 (d, *J*=2.8 Hz, 1H), 3.31 (s, 1H), 2.87 (dd, *J* = 13.2; 4.2 Hz, 2H), 1.93 (m, 1H), 1.65 (m, 1H), 1.44 (s, 3H), 1.21 (s, 3H), 1.02 (m, 1H), 0.47 (m, 3H), 0.36 (m, 1H) ppm. ES MS *M*+1 = 373.

EXAMPLE 9

5-(Dimethylaminocarbonyl)-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione



Step 1: 2-(4-Fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione-5-carboxylic acid.


A solution of 5-(tert-butyloxycarbonyl)-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione, Diastereomer 1 from Example 4, (0.015 g, 0.037 mmol) in 50 % TFA in CH₂Cl₂ (2.4 mL) was stirred at room temperature. After 1.3 hours, TFA acid




(0.5 mL) was added to the reaction mixture. After 48 hours, the reaction mixture was concentrated under vacuum to give the title compound. ES MS $M+1 = 349$.




Step 2: 5-(Dimethylaminocarbonyl)-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione.

To a solution of 2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione-5-carboxylic acid (0.012 g, 0.0034 mol) in DMF (0.5 mL) was added EDC (0.007 g, 0.038 mmol), dimethylamine hydrochloride (0.003 g, 0.038 mmol) and HOBT (0.006 g, 0.038 mmol). After stirring at room temperature, triethylamine (0.008 g, 0.076 mmol) was added. After 2 hours, the reaction mixture was concentrated under vacuum. The residual material was partitioned between CHCl_3 and water. The organic extract was dried with Na_2SO_4 , filtered, and concentrated under vacuum. The residue was purified using reverse phase HPLC on a C18 stationary phase eluting with 15 % -85 % acetonitrile (0.1 % TFA) in H_2O (0.1 % TFA) to provide the title compound. ^1H NMR (400 MHz, CDCl_3) δ 7.25 (m, 2H), 7.04 (t, $J=8.6$ Hz, 2H), 4.75 (m, 1H), 4.44 (m, 1H), 3.95 (d, $J=12.3$ Hz, 1H), 3.29 (m, 4H), 3.18 (s, 3H), 1.81 (m, 1H), 1.65 (s, 6H), 1.49 (m, 1H) ppm. ES MS $M+1 = 376$.

EXAMPLES 10 TO 16

Example	Compound	Data
10	<p>2-(3-Chloro-4-fluorobenzyl)-8-hydroxy-5,5,6-trimethyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione</p>  <p>The title compound was prepared using procedures analogous to Steps 1-3 of Example 1, with 3-chloro-4-fluorobenzyl bromide replacing 4-fluorobenzyl bromide, and then using procedures analogous to Steps 1-4 of Example 2.</p> <p>The enantiomers of the title compound were separated with a ChiralPak AD column with 100% methanol as the mobile phase, wherein the first enantiomer had a retention time of 5.58 min and a negative sign of rotation, and the second enantiomer had a retention time of 7.67 min and a positive sign of rotation.</p>	<p>^1H NMR (400 MHz, CDCl_3) δ 7.32 (m, 1H), 7.16 (m, 1H), 7.12 (t, $J=8$ Hz, 1H), 4.64 (d, $J=16$ Hz, 1H), 4.52 (d, $J=16$ Hz, 1H), 3.34 (m, 2H), 3.02 (s, 3H), 2.84 (dd, $J = 13.2$; 4.2 Hz, 1H), 2.46 (br, 1H), 1.91 (m, 1H), 1.65 (m, 1H), 1.38 (s, 3H), 1.14 (s, 3H) ppm. ES MS $M+1 = 367$.</p>

11	<p>6'-(4-Fluorobenzyl)-4'-hydroxy-2'-methyl-6',7',8',8a'-tetrahydro-2' <i>H</i>-spiro[cyclopentane-1,1'-[2,6]naphthyridine]-3',5'-dione</p>  <p>The title compound was prepared using procedures analogous to Example 2 except that nitrocyclopentane was used in Step 1 instead of 2-nitropropane.</p> <p>The enantiomers of the title compound were separated using a ChiralPak AD column with 50% methanol in ethanol as the mobile phase, wherein the first enantiomer had a retention time of 5.05 min and a negative sign of rotation, and the second enantiomer had a retention time of 5.67 min and a positive sign of rotation.</p>	<p>¹H NMR (400 MHz, CDCl₃) δ 7.26 (m, 2H), 7.04 (t, <i>J</i>=8.7 Hz, 2H), 4.67 (d, <i>J</i>=14.5 Hz, 1H), 4.52 (d, <i>J</i>=14.6 Hz, 1H), 3.29 (m, 2H), 3.03 (s, 3H), 2.93 (dd, <i>J</i>=3.84; 13.2 Hz, 1H), 2.22 (m, 3H), 1.96 (m, 1H), 1.82 (m, 1H), 1.67 (m, 2H), 1.45 (m, 1H), 1.31 (m, 1H) ppm. ES MS <i>M</i>+1 = 359.</p>
12	<p>2-(3,4-difluorobenzyl)-8-hydroxy-5,5,6-trimethyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione</p>  <p>The title compound was prepared using procedures analogous to Steps 1-3 of Example 1, with 3,4-difluorobenzyl bromide replacing 4-fluorobenzyl bromide, and then using procedures analogous to Steps 1-4 of Example 2.</p>	<p>¹H NMR (400 MHz, CDCl₃) δ 7.13 (m, 2H), 7.02 (m, 1H), 4.66 (d, <i>J</i>=16 Hz, 1H), 4.49 (d, <i>J</i>=16 Hz, 1H), 3.34 (m, 2H), 3.02 (s, 3H), 2.84 (dd, <i>J</i> = 13, 4 Hz, 1H), 1.93 (m, 1H), 1.67 (m, 1H), 1.39 (s, 3H), 1.13 (s, 3H) ppm. ES MS <i>M</i>+1 = 351.</p>
13	<p>6'-(4-Fluorobenzyl)-4'-hydroxy-2'-methyl-6',7',8',8a'-tetrahydro-2' <i>H</i>-spiro[cyclobutane-1,1'-[2,6]naphthyridine]-3',5'-dione</p>  <p>The title compound was prepared via Step 1 of Example 2 using nitrocyclobutane in place of 2-nitropropane, followed by Step 3 of Example 5, and then Steps 1 and 2 of Example 7.</p>	<p>¹H NMR (400 MHz, CDCl₃) δ 7.27 (m, 2H), 7.05 (t, <i>J</i>=8.4 Hz, 2H), 4.69 (d, <i>J</i>=14.5 Hz, 1H), 4.54 (d, <i>J</i>=14.7 Hz, 1H), 3.41 (m, 2H), 3.21 (s, 3H), 2.84 (m, 2H), 2.34 (m, 1H), 2.22 (m, 1H), 2.07 (m, 2H), 1.94 (m, 2H), 1.67 (m, 1H) ppm. ES MS <i>M</i>+1 = 345.</p>

14	<p>5-[(2-Methylpropyl)aminocarbonyl]-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione</p>  <p>The title compound was prepared via a procedure analogous to Step 2 of Example 9, using the appropriate amine in place of dimethylamine hydrochloride.</p>	<p>¹H NMR (400 MHz, CD₃OD) δ 7.33 (m, 2H), 7.07 (t, <i>J</i>=8.8 Hz, 2H), 4.73 (d, <i>J</i>=14.6 Hz, 1H), 4.51 (d, <i>J</i>=14.6 Hz, 1H), 3.92 (d, <i>J</i>=12.4 Hz, 1H), 3.40 (m, 1H), 3.11 (m, 4H), 2.93 (s, 3H), 1.83 (m, 2H), 1.61 (m, 1H), 0.94 (s, 6H) ppm. ES MS <i>M</i>+1 = 404.</p>
15	<p>5-(tert-Butylaminocarbonyl)-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione</p>  <p>The title compound was prepared via a procedure analogous to Step 2 of Example 9, using the appropriate amine in place of dimethylamine hydrochloride.</p>	<p>ES MS <i>M</i>+1 = 404.</p>
16	<p>5-[(2-Pyridylmethyl)aminocarbonyl]-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione</p> 	<p>ES MS <i>M</i>+1 = 439.</p>

EXAMPLE 17

Oral Compositions

- 5 As a specific embodiment of an oral composition of a compound of this invention, 50 mg of compound of Example 1 is formulated with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size 0 hard gelatin capsule. Encapsulated oral compositions containing any one of the compounds of Examples 2-16 can be similarly prepared.

EXAMPLE 18

HIV Integrase Assay: Strand Transfer Catalyzed by Recombinant Integrase

Assays for the strand transfer activity of integrase were conducted in accordance with WO 02/30930 for recombinant integrase. Representative compounds of the present invention exhibit inhibition of strand transfer activity in this assay. For example, the compounds prepared in Examples 1-16 were tested in the integrase assay and all were found to have IC₅₀'s less than about 1 micromolar.

Further description on conducting the assay using preassembled complexes is found in . Wolfe, A.L. et al., *J. Virol.* 1996, 70: 1424-1432, Hazuda et al., *J. Virol.* 1997, 71: 7005-7011; Hazuda et al., *Drug Design and Discovery* 1997, 15: 17-24; and Hazuda et al., *Science* 2000, 287: 646-650.

EXAMPLE 19


Assay for inhibition of HIV replication

Assays for the inhibition of acute HIV infection of T-lymphoid cells were conducted in accordance with Vacca, J.P. et al., *Proc. Natl. Acad. Sci. USA* 1994, 91: 4096. Representative compounds of the present invention exhibit inhibition of HIV replication in this assay. For example, the compounds prepared in Examples 1-16 were found to have IC₉₅'s of less than about 10 micromolar in the present assay.

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, the practice of the invention encompasses all of the usual variations, adaptations and/or modifications that come within the scope of the following claims.

WHAT IS CLAIMED IS:

1. A compound of Formula I, or a pharmaceutically acceptable salt thereof:



5 wherein:

R¹ is -C₁₋₆ alkyl substituted with R^J, wherein R^J is:

- (A) aryl or aryl fused to a 5- or 6-membered heteroaromatic ring containing from 1 to 4 heteroatoms independently selected from N, O and S, wherein the aryl or fused aryl is

10 (i) optionally substituted with from 1 to 5 substituents each of which is independently:

- (1) -C₁₋₆ alkyl, which is optionally substituted with -OH, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, -CN, -NO₂, -N(R^A)R^B, -C(=O)N(R^A)R^B, -C(=O)R^A, -CO₂R^A, -S(O)_nR^A, -SO₂N(R^A)R^B, -N(R^A)C(=O)R^B, -N(R^A)CO₂R^B, -N(R^A)SO₂R^B, -N(R^A)SO₂N(R^A)R^B, -OC(=O)N(R^A)R^B, or -N(R^A)C(=O)N(R^A)R^B,

- (2) -O-C₁₋₆ alkyl,
- (3) -C₁₋₆ haloalkyl,
- (4) -O-C₁₋₆ haloalkyl,

- (5) -OH,
- (6) halo,
- (7) -CN,
- (8) -NO₂,
- (9) -N(RA)RB,

- $$\begin{aligned} (10) \quad & -\text{C}(=\text{O})\text{N}(\text{R}^{\text{A}})\text{R}^{\text{B}}, \\ (11) \quad & -\text{C}(=\text{O})\text{R}^{\text{A}}, \\ (12) \quad & -\text{CO}_2\text{R}^{\text{A}}, \\ (13) \quad & -\text{SR}^{\text{A}}, \\ (14) \quad & -\text{S}(=\text{O})\text{R}^{\text{A}}, \end{aligned}$$

- (15) $-\text{SO}_2\text{RA}$,
 (16) $-\text{SO}_2\text{N(RA)RB}$,
 (17) $-\text{N(RA)SO}_2\text{RB}$,
 (18) $-\text{N(RA)SO}_2\text{N(RA)RB}$,
 5 (19) $-\text{N(RA)C(=O)RB}$,
 (20) $-\text{N(RA)C(=O)-C(=O)N(RA)RB}$, or
 (21) $-\text{N(RA)CO}_2\text{RB}$, and
- (ii) optionally substituted with 1 or 2 substituents each of which is independently:
- 10 (1) aryl,
 (2) $-\text{C}_{1-6}$ alkyl substituted with aryl,
 (3) $-\text{HetA}$,
 (4) $-\text{C(=O)-HetA}$; or
 (5) $-\text{HetB}$;
- 15 wherein each HetA is independently a C_{4-7} azacycloalkyl or a C_{3-6} diazacycloalkyl, either of which is optionally substituted with from 1 to 3 substituents each of which is independently oxo or C_{1-6} alkyl; and
- 20 wherein each HetB is a 5- or 6-membered heteroaromatic ring containing from 1 to 4 heteroatoms independently selected from N, O and S, wherein the heteroaromatic ring is optionally substituted with from 1 to 4 substituents each of which is independently halo, $-\text{C}_{1-6}$ alkyl, $-\text{C}_{1-6}$ haloalkyl, $-\text{O-C}_{1-6}$ alkyl, $-\text{O-C}_{1-6}$ haloalkyl, or hydroxy; or
- (B) a 5- or 6-membered heteroaromatic ring containing from 1 to 4 heteroatoms independently selected from N, O and S; wherein the heteroaromatic ring is:
- 25 (i) optionally substituted with from 1 to 4 substituents each of which is independently halogen, $-\text{C}_{1-6}$ alkyl, $-\text{C}_{1-6}$ haloalkyl, $-\text{O-C}_{1-6}$ alkyl, $-\text{O-C}_{1-6}$ haloalkyl, or hydroxy, and
- (ii) optionally substituted with 1 or 2 substituents each of which is independently aryl or $-\text{C}_{1-6}$ alkyl substituted with aryl;
- 30 R^2 , R^3 , R^4 and R^5 are defined as follows:
- (A) R^2 , R^3 , R^4 and R^5 are each independently:
- (1) $-\text{H}$,
 (2) $-\text{C}_{1-6}$ alkyl, which is optionally substituted with $-\text{OH}$, $-\text{O-C}_{1-6}$ alkyl, $-\text{O-C}_{1-6}$ haloalkyl, $-\text{CN}$, $-\text{N(RA)RB}$, $-\text{C(=O)N(RA)RB}$, $-\text{C(=O)RA}$, $-\text{CO}_2\text{RA}$, $-\text{S(O)}_n\text{RA}$,

-SO₂N(RA)RB, -N(RA)C(=O)RB, -N(RA)CO₂RB, -N(RA)SO₂RB,
-N(RA)SO₂N(RA)RB, -N(RA)C(=O)N(RA)RB, or -OC(=O)N(RA)RB,

- (3) -C₁₋₆ haloalkyl,
- (4) CycA,
- (5) AryA,
- (6) HetC, or
- (7) -C₁₋₆ alkyl substituted with CycA, AryA, or HetC;

(B) R² and R⁴ together with the carbon atoms to which each is attached form a carbon-carbon double bond; and R³ and R⁵ are each independently as defined in part A above;

(C) R² and R³ together with the carbon atom to which they are both attached form a 3- to 8-membered saturated carbocyclic ring which is optionally substituted with from 1 to 4 substituents each of which is independently -OH, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, or -O-C₁₋₆ haloalkyl; and R⁴ and R⁵ are each independently as defined in part A above; or

(D) R⁴ and R⁵ together with the carbon atom to which they are both attached form a 3- to 8-membered saturated carbocyclic ring which is optionally substituted with from 1 to 4 substituents each of which is independently -OH, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, or -O-C₁₋₆ haloalkyl; and R² and R³ are each independently as defined in part A above;

R⁶ is:

- (1) -H,
- (2) -C₁₋₆ alkyl, which is optionally substituted with -OH, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, -CN, -N(RA)RB, -C(=O)N(RA)RB, -C(=O)RA, -CO₂RA, -S(O)_nRA, -SO₂N(RA)RB, -N(RA)C(=O)RB, -N(RA)CO₂RB, -N(RA)SO₂RB, -N(RA)SO₂N(RA)RB, -N(RA)C(=O)N(RA)RB, or -OC(=O)N(RA)RB,
- (3) -C₁₋₆ haloalkyl,
- (4) CycA,
- (5) AryA,
- (6) HetC, or
- (7) -C₁₋₆ alkyl substituted with CycA, AryA, or HetC;

R⁷ and R⁸ are each independently:

- (1) -H,

- (2) -C₁₋₆ alkyl, which is optionally substituted with -OH, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, -CN, -N(RA)R^B, -C(=O)N(RA)R^B, -C(=O)RA, -CO₂RA, -S(O)_nRA, -SO₂N(RA)R^B, -N(RA)C(=O)R^B, -N(RA)CO₂R^B, -N(RA)SO₂R^B, -N(RA)SO₂N(RA)R^B, -N(RA)C(=O)N(RA)R^B, or -OC(=O)N(RA)R^B,
 5 (3) -C₁₋₆ haloalkyl,
 (4) -C(=O)RA,
 (5) -CO₂RA,
 (6) -C(=O)N(RA)R^B,
 (7) -N(RA)SO₂N(RA)R^B,
 10 (8) -R^K,
 (9) -C(=O)-R^K,
 (10) -C(=O)N(RA)-R^K,
 (11) -C(=O)N(RA)-C₁₋₆ alkylene-R^K, or
 (12) -C₁₋₆ alkyl substituted with -R^K, -C(=O)-R^K, -C(=O)N(RA)-R^K, or -C(=O)N(RA)-C₁₋₆
 15 alkylene-R^K;

or alternatively R⁷ and R⁸ together with the carbon atom to which they are both attached form a 3- to 8-membered saturated carbocyclic ring which is optionally substituted with from 1 to 4 substituents each of which is independently halogen, -OH, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, or -O-C₁₋₆ haloalkyl;

20

R⁹ is:

- (1) -H,
 (2) -C₁₋₆ alkyl, which is optionally substituted with -OH, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, -CN, -N(RA)R^B, -C(=O)N(RA)R^B, -C(=O)RA, -CO₂RA, -S(O)_nRA, -SO₂N(RA)R^B,
 25 -N(RA)C(=O)R^B, -N(RA)CO₂R^B, -N(RA)SO₂R^B, -N(RA)SO₂N(RA)R^B,
 -N(RA)C(=O)N(RA)R^B, or -OC(=O)N(RA)R^B,
 (3) -C₁₋₆ haloalkyl,
 (4) CycA,
 (5) AryA,
 30 (6) HetC, or
 (7) -C₁₋₆ alkyl substituted with CycA, AryA, or HetC;

each n is independently an integer equal to zero, 1, or 2;

each R^A is independently H or C₁₋₆ alkyl;

each R^B is independently H or C₁₋₆ alkyl;

5 each R^K is independently CycA, AryA, or HetC;

each CycA is independently a C₃₋₈ cycloalkyl, which is optionally substituted with from 1 to 4 substituents each of which is halogen, -OH, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, or -O-C₁₋₆ haloalkyl;

10

each AryA is independently an aryl, which is

- (a) optionally substituted with from 1 to 5 substituents each of which is independently -C₁₋₆ alkyl, -C₁₋₆ alkylene-OH, -C₁₋₆ alkylene-O-C₁₋₆ alkyl, -C₁₋₆ alkylene-O-C₁₋₆ haloalkyl, -C₁₋₆ alkylene-N(R^A)R^B, -C₁₋₆ alkylene-C(=O)N(R^A)R^B, -C₁₋₆ alkylene-C(=O)R^A, -C₁₋₆ alkylene-CO₂R^A, -C₁₋₆ alkylene-S(O)_nR^A, -O-C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ haloalkyl, -OH, halo, -N(R^A)R^B, -C(=O)N(R^A)R^B, -C(=O)R^A, -CO₂R^A, -S(O)_nR^A, or -SO₂N(R^A)R^B, and
- (b) optionally substituted with C₃₋₈ cycloalkyl, aryl, HetD, or -C₁₋₆ alkyl substituted with C₃₋₈ cycloalkyl, aryl, or HetD;

20

each HetC is independently a 4- to 7-membered saturated or unsaturated heterocyclic ring containing at least one carbon atom and from 1 to 4 heteroatoms independently selected from N, O and S, wherein the heterocyclic ring is

- (a) optionally substituted with from 1 to 4 substituents each of which is halogen, -C₁₋₆ alkyl, -C₁₋₆ haloalkyl, -O-C₁₋₆ alkyl, -O-C₁₋₆ haloalkyl, OH, or oxo, and
- (b) optionally substituted with C₃₋₈ cycloalkyl, aryl, HetD, or -C₁₋₆ alkyl substituted with C₃₋₈ cycloalkyl, aryl, or HetD;

25

each HetD is independently a 4- to 7-membered saturated or unsaturated heterocyclic ring containing at least one carbon atom and from 1 to 4 heteroatoms independently selected from N, O and S; and

30

each aryl is independently (i) phenyl or (ii) a 9- or 10-membered bicyclic, fused carbocyclic ring system in which at least one ring is aromatic.

2. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein R^1 is $-CH_2-R^J$, and R^J is phenyl, pyridyl, quinolinyl, isoquinolinyl, cinnolinyl, or quinazolinyl, any of which is

(a) optionally substituted with from 1 to 4 substituents each of which is independently:

- (1) $-C_{1-4}$ alkyl,
- (2) $-O-C_{1-4}$ alkyl,
- (3) $-C_{1-4}$ haloalkyl,
- (4) $-O-C_{1-4}$ haloalkyl,
- (5) halo,
- (6) $-CN$,
- (7) $-N(R^A)R^B$,
- (8) $-C(=O)N(R^A)R^B$,
- (9) $-S(=O)R^A$,
- (10) $-SO_2R^A$,
- (11) $-N(R^A)SO_2R^B$,
- (12) $-N(R^A)SO_2N(R^A)R^B$,
- (13) $-N(R^A)C(=O)R^B$, or
- (14) $-N(R^A)C(=O)-C(=O)N(R^A)R^B$, and

(b) optionally substituted with phenyl, benzyl, $-HetA$, or $-C(=O)-HetA$; wherein each $HetA$ is independently a C_{4-7} azacycloalkyl or a C_{3-6} diazacycloalkyl, either of which is optionally substituted with from 1 to 3 substituents each of which is independently oxo or C_{1-4} alkyl; and with the proviso that when $HetA$ is attached to the rest of the compound via the $-C(=O)-$ moiety, the $HetA$ is attached to the $-C(=O)-$ via a ring N atom.

3. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein R^2 , R^3 , R^4 and R^5 are defined as follows:

- (A) R^2 and R^4 are as defined in part A of claim 1, and R^3 and R^5 are both H;
- (B) R^2 and R^4 are as defined in part B of claim 1; and R^3 and R^5 are both H;
- (C) R^2 and R^3 are as defined in part C of claim 1; and R^4 and R^5 are both H; or
- (D) R^4 and R^5 are as defined in part D of claim 1; and R^2 and R^3 are both H.

4. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein R^6 is:

- (1) $-H$,

- (2) -C₁₋₆ alkyl,
 (3) -C₁₋₆ fluoroalkyl,
 (4) CycA,
 (5) AryA, or
 5 (6) -C₁₋₆ alkyl substituted with AryA;

5. The compound according to claim 4, or a pharmaceutically acceptable salt thereof, wherein R⁶ is H.

10 6. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein R⁷ and R⁸ are each independently:


- (1) -H,
 (2) -C₁₋₆ alkyl,
 (3) -CO₂RA,
 15 (4) -C(=O)N(RA)RB,
 (5) -RK,
 (6) -C(=O)-RK,
 (7) -C(=O)N(RA)-RK, or
 (8) -C(=O)N(RA)-C₁₋₆ alkylene-RK;

20 or alternatively R⁷ and R⁸ together with the carbon atom to which they are both attached form a 3- to 7-membered saturated carbocyclic ring.

25 7. The compound according to claim 1, or a pharmaceutically acceptable salt thereof, wherein R⁹ is:

- (1) -H,
 (2) -C₁₋₆ alkyl
 (3) -C₁₋₆ fluoroalkyl,
 (4) CycA, or
 30 (5) -C₁₋₆ alkyl substituted with CycA, AryA, or HetC.

8. A compound of Formula II, or a pharmaceutically acceptable salt thereof:



wherein:

- 5 X¹ and X² are each independently -H, -C₁₋₄ alkyl, -O-C₁₋₄ alkyl, -C₁₋₄ haloalkyl, -O-C₁₋₄ haloalkyl, halo, -CN, -N(R^A)R^B, -C(=O)N(R^A)R^B, or -S(O)_nR^A, wherein n is an integer equal to zero, 1, or 2;

R², R³, R⁴ and R⁵ are defined as follows:

- 10 (A) R² and R⁴ are each independently -H, -C₁₋₄ alkyl, -C₁₋₄ fluoroalkyl, C₃₋₆ cycloalkyl, phenyl, or benzyl; and R⁴ and R⁵ are both H;
- (B) R² and R⁴ together with the carbon atoms to which each is attached form a carbon-carbon double bond; and R³ and R⁵ are both H;
- (C) R² and R³ together with the carbon atom to which they are both attached form cyclopropyl; and R⁴ and R⁵ are both H; or
- 15 (D) R⁴ and R⁵ together with the carbon atom to which they are both attached form cyclopropyl; and R² and R³ are both H;

R⁶ is H, -C₁₋₄ alkyl, CF₃, cyclopropyl, phenyl or benzyl;

- 20 R⁷ is H or -C₁₋₄ alkyl;

R⁸ is -H, -C₁₋₄ alkyl, -CO₂-C₁₋₄ alkyl, -C(=O)NH(C₁₋₄ alkyl), -C(=O)N(C₁₋₄ alkyl)₂, C₃₋₆ cycloalkyl, -C(=O)-HetE, or -C(=O)N(R^A)-(CH₂)₁₋₂-HetF; wherein

- 25 HetE is a 4- to 7-membered saturated heterocyclic ring containing at least one carbon atom and from 1 to 4 heteroatoms selected from 1 to 4 N atoms, zero or 1 oxygen atoms, and zero or 1 sulfur atoms, wherein the saturated heterocyclic is optionally substituted with from 1 to 3 substituents each of which is independently oxo or C₁₋₄ alkyl; and with the proviso that the saturated heterocyclic is attached to the -C(=O)- via a ring N atom; and

HetF is a 5- or 6-membered heteroaromatic ring containing from 1 to 4 heteroatoms independently selected from N, O and S, wherein the heteroaromatic ring is optionally substituted with 1 or 2 substituents each of which is independently a C₁₋₄ alkyl;

5 or alternatively R⁷ and R⁸ together with the carbon atom to which they are both attached form a 3- to 6-membered saturated carbocyclic ring;

R⁹ is -H, -C₁₋₄ alkyl, -CH₂CF₃, -C₃₋₆ cycloalkyl, -CH₂-C₃₋₆ cycloalkyl, or -CH₂-phenyl;

10 each R^A is independently H or C₁₋₄ alkyl; and

each R^B is independently H or C₁₋₄ alkyl.

15 9. A compound according to claim 8, or a pharmaceutically acceptable salt thereof, wherein:

X¹ and X² are each independently H, fluoro, chloro, methyl, trifluoromethyl, methoxy, CN, -SO₂CH₃, -C(=O)NH(CH₃), or -C(=O)N(CH₃)₂;


20 R², R³, R⁴ and R⁵ are all H;


R⁶ is H, methyl, cyclopropyl, or phenyl;

R⁷ is H or methyl;

25 R⁸ is -H, -C₁₋₄ alkyl, -CO₂-C₁₋₄ alkyl, -C(=O)NH(C₁₋₄ alkyl), -C(=O)N(C₁₋₄ alkyl)₂, C₃₋₆ cycloalkyl, -C(=O)-HetE, or -C(=O)N(R^A)-(CH₂)₁₋₂-HetF; wherein

HetE is selected from the group consisting of:



30 , and wherein the asterisk * denotes the point of attachment to the -C(=O) moiety; and

HetF is selected from the group consisting of pyrrolyl, imidazolyl, triazolyl, tetrazolyl, oxazolyl, isooxazolyl, pyridyl, pyrimidinyl, and pyrazinyl;

or alternatively R⁷ and R⁸ together with the carbon atom to which they are both attached form cyclopropyl, cyclobutyl, cyclopentyl, or cyclohexyl; and

R⁹ is H, methyl, ethyl, n-propyl, isopropyl, -CH₂CF₃, cyclopropyl, or -CH₂-cyclopropyl.

10. A compound, or a pharmaceutically acceptable salt thereof, selected from the group consisting of:

2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione;

2-(4-fluorobenzyl)-8-hydroxy-5,5,6-trimethyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione;

2-(4-fluorobenzyl)-8-hydroxy-6-methyl-4a-phenyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione;

5-(tert-butyloxycarbonyl)-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione;

5-ethyl-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione;

(4aR, 5R)-5-ethyl-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione;

(4aR, 5S)-5-ethyl-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione;

6-(cyclopropylmethyl)-2-(4-fluorobenzyl)-8-hydroxy-5,5-dimethyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione;

5-(dimethylaminocarbonyl)-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione;

2-(3-chloro-4-fluorobenzyl)-8-hydroxy-5,5,6-trimethyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione;

5 6'-(4-fluorobenzyl)-4'-hydroxy-2'-methyl-6',7',8',8a'-tetrahydro-2' *H*-spiro[cyclopentane-1,1'-[2,6]naphthyridine]-3',5'-dione;

2-(3,4-difluorobenzyl)-8-hydroxy-5,5,6-trimethyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione;

10 6'-(4-fluorobenzyl)-4'-hydroxy-2'-methyl-6',7',8',8a'-tetrahydro-2' *H*-spiro[cyclobutane-1,1'-[2,6]naphthyridine]-3',5'-dione;

5-[(2-methylpropyl)aminocarbonyl]-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione;

15 5-(tert-butylaminocarbonyl)-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione; and

20 5-[(2-pyridylmethyl)aminocarbonyl]-2-(4-fluorobenzyl)-8-hydroxy-6-methyl-2,3,4,4a,5,6-hexahydro-2,6-naphthyridine-1,7-dione.

11. A pharmaceutical composition comprising an effective amount of a compound according to any one of claims 1 to 10, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

25 12. A method of inhibiting HIV integrase in a subject in need thereof which comprises administering to the subject an effective amount of the compound according to any one of claims 1 to 10, or a pharmaceutically acceptable salt thereof.

30 13. A method for preventing or treating infection by HIV or for preventing, treating or delaying the onset of AIDS in a subject in need thereof which comprises administering to the subject an effective amount of the compound according to any one of claims 1 to 10, or a pharmaceutically acceptable salt thereof.

14. Use of a compound according to any one of claims 1 to 10, or a pharmaceutically acceptable salt thereof, for inhibiting HIV integrase in a subject in need thereof.

5 15. Use of a compound according to any one of claims 1 to 10, or a pharmaceutically acceptable salt thereof, for preventing or treating infection by HIV or for preventing, treating or delaying the onset of AIDS in a subject in need thereof.


10 16. A compound according to any one of claims 1 to 10, or a pharmaceutically acceptable salt thereof, for use in the preparation of a medicament for inhibiting HIV integrase in a subject in need thereof.

15 17. A compound according to any one of claims 1 to 10, or a pharmaceutically acceptable salt thereof, for use in the preparation of a medicament for preventing or treating infection by HIV or for preventing, treating or delaying the onset of AIDS in a subject in need thereof.

20 18. A pharmaceutical combination which is (i) a compound according to any one of claims 1 to 10, or a pharmaceutically acceptable salt thereof, and (ii) an HIV infection/AIDS antiviral agent selected from the group consisting of HIV protease inhibitors, non-nucleoside HIV reverse transcriptase inhibitors and nucleoside HIV reverse transcriptase inhibitors; wherein the compound of (i) or its pharmaceutically acceptable salt and the HIV infection/AIDS antiviral agent of (ii) are each employed in an amount that renders the combination effective for inhibiting HIV integrase, for preventing or treating infection by HIV, or for preventing, treating or delaying the onset of AIDS.

ABSTRACT OF THE DISCLOSURE

Hydroxy (tetra- or hexa-)hydronaphthyridine dione compounds of Formula I are inhibitors of HIV integrase and inhibitors of HIV replication:



- 5 wherein R¹, R², R³, R⁴, R⁵, R⁶, R⁷, R⁸ and R⁹ are defined herein. The compounds are useful in the prevention and treatment of infection by HIV and in the prevention, delay in the onset, and treatment of AIDS. The compounds are employed against HIV infection and AIDS as compounds per se or in the form of pharmaceutically acceptable salts. The compounds and their salts can be employed as ingredients in pharmaceutical compositions, optionally in combination with other antivirals,
- 10 immunomodulators, antibiotics or vaccines.